



Generalized Elliptic Integrals Involving Incomplete Fox Wright Function

Shalini Shekhawat¹, Akanksha Shukla^{2,*}, Kanak Modi²

¹Department of Mathematics, SKIT M&G, Jaipur 302017, India

²Department of Mathematics, Amity University of Rajasthan, Jaipur 302 015, India

Received 24 August 2021; Received in revised form 20 November 2021

Accepted 27 November 2021; Available online 30 December 2021

ABSTRACT

Elliptic integrals are used in various diverse physical fields (like some exclusive radiation field problems, analysis of crystallographic minimum surface problem, study of electromagnetic waves through an elliptic disk, elliptic crack problems etc.). Due to these applications a remarkably large number of elliptic integrals are defined and studied earlier. Motivated by these works we have derived some new elliptic integrals involving incomplete wright function. The results established in this paper are basic in nature and can be reduced in many special cases. Some of the special cases are mentioned here as corollaries.

Keywords: Beta and Gamma functions; Fox wright function; Incomplete elliptic integral; Riemann-Liouville fractional differential operator

1. Introduction

Nowadays, Research is centered on integral presentation of different type of special functions due to their use of various fields. Several integral formulas having generalized special function, plays an important role in physical and technical problems. The generalized special function with contrasting line of research have inspired the researchers to look into the area of integrals and linked generalized special function [1]. With the use of Riemann-Liouville fractional integral we describe many different hypergeometric notations

and then we can use these representations for the derivation of many definite integrals [2-4]. Incomplete elliptic integrals and related definite integrals help to solve many physical problems and may find applications in definite engineering problems. Inspired by this idea, in this paper we estimated certain incomplete integrals and related general definite integrals. Here our target is to provide the theory of generalized incomplete elliptic integrals in an exclusive and generalized way.

Incomplete Fox-Wright Ψ -functions

${}_r\psi_s^{(\Gamma)}(x)$ and ${}_r\psi_s^{(\gamma)}(x)$:

Here we define the family of the incomplete Fox-Wright function ${}_r\psi_s^{(\Gamma)}(x)$ and ${}_r\psi_s^{(\gamma)}(x)$, in both series and contour integral form which is a generalization of incomplete hypergeometric functions defined by Srivastava et al. [9, 13].

$${}_r\psi_s^{(\Gamma)} \left[\begin{matrix} (a_1, \alpha_1, x), (a_j, \alpha_j)_{2,r} \\ (b_j, \beta_j)_{1,s} \end{matrix} \middle| x \right] = \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \frac{x^n}{n!}, \quad (1.1)$$

$$= \frac{1}{2\pi i} \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)^c} \int_C \frac{\Gamma(a_1 + v, y) \prod_{j=2}^r \Gamma(a_j + v)}{\prod_{j=1}^s \Gamma(b_j + v)} \Gamma(-v)(-x)^v dv \left(|\arg(-x)| < \pi \right), \quad (1.2)$$

and

$$(\alpha_j > 0 \quad (j=1, \dots, r); \beta_j > 0 \quad (j=1, \dots, s); 1 + \sum_{j=1}^s \beta_j - \sum_{j=1}^r \alpha_j \geq 0).$$

Similarly, Eq. (1.3) represents the series representation and Eq. (1.4) represents the contour integral of incomplete Fox-Wright function ${}_r\psi_s^{(\gamma)}(x)$:

$${}_r\psi_s^{(\gamma)} \left[\begin{matrix} (a_1, \alpha_1, x), (a_j, \alpha_j)_{2,r} \\ (b_j, \beta_j)_{1,s} \end{matrix} \middle| x \right] = \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \frac{x^n}{n!}, \quad (1.3)$$

$$= \frac{1}{2\pi i} \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)^c} \int_C \frac{\gamma(a_1 + v, y) \prod_{j=2}^r \Gamma(a_j + v)}{\prod_{j=1}^s \Gamma(b_j + v)} \Gamma(-v)(-x)^v dv \left(|\arg(-x)| < \pi \right) \quad (1.4)$$

where

$$(\alpha_j > 0 \quad (j=1, \dots, r); \beta_j > 0 \quad (j=1, \dots, s);$$

$$1 + \sum_{j=1}^s \beta_j - \sum_{j=1}^r \alpha_j \geq 0).$$

$$\alpha_j > 0 \quad (j=1, \dots, r); \beta_j > 0 \quad (j=1, \dots, s);$$

$$1 + \sum_{j=1}^s \beta_j - \sum_{j=1}^r \alpha_j \geq 0).$$

The multivariable hypergeometric function defined by Srivastava & Daoust [5-7] is given by

$$F_{n, l_1, l_2, \dots, l_k}^{r, s_1, s_2, \dots, s_k} \left[\begin{matrix} (a_j; \alpha_j', \dots, \alpha_j^{(k)})_{1,r} : (c_j', t_j')_{1, s_1}; \dots; \\ (b_j; \beta_j', \dots, \beta_j^{(k)})_{1,n} : (d_j', \delta_j')_{1, l_1}; \dots; \\ (c_j^{(k)}, t_j^{(k)})_{1, s_k} \\ (d_j^{(r)}, \delta_j^{(r)})_{1, l_k} \end{matrix} \middle| x_1, x_2, \dots, x_k \right] = \sum_{m_1, \dots, m_k}^{\infty} \frac{\prod_{j=1}^r (a_j)_{m_1 \alpha_j' + \dots + m_k \alpha_j^{(k)}}}{\prod_{j=1}^n (b_j)_{m_1 \beta_j' + \dots + m_k \beta_j^{(k)}}} \frac{\prod_{j=1}^{s_1} (c_j')_{m_1 t_j'} \dots \prod_{j=1}^{s_k} (c_j^{(k)})_{m_1 t_j^{(k)}}}{\prod_{j=1}^{m_1} (d_j')_{m_1 \delta_j'} \dots \prod_{j=1}^{m_r} (d_j^{(k)})_{m_1 \delta_j^{(k)}}} \quad (5)$$

with variable and parametric constraints, the above-mentioned series is absolutely convergent. The Pochhammer symbol denoted by $(c)_p$ is defined

as:

$$(c)_p = \frac{\Gamma(c+p)}{\Gamma(c)} = \begin{cases} 1, & \{p=0; \lambda \in C / \{0\}\} \\ c(c+1) \dots (c+n-1), & \{p=n \in N = N_0; c \in C\} \end{cases}$$

where $c, p \in C$. (1.6)

Riemann-Liouville operator $D_z^\delta f(z)$ of fractional calculus [8]:

$$D_Z^{\gamma-\delta} \left\{ z^{\gamma-1} \prod_{j=1}^k \left\{ \left(1 - a_j z^{\delta_j} \right)^{-\alpha_j} \right\} \right\} \\ = \frac{\Gamma(\gamma)}{\Gamma(\delta)} z^{\delta-1} F_{1:0;\dots;0}^{1:1;\dots;1} \left[\begin{matrix} (\gamma; \delta_1, \dots, \delta_k) : (\alpha_1, 1); \\ (\delta; \delta_1, \dots, \delta_k); \dots; \\ \dots; (\alpha_k, 1) \end{matrix} \middle| \begin{matrix} a_1 z^{\delta_1}, \dots, a_k z^{\delta_k} \end{matrix} \right] \quad (1.7)$$

where $R(\gamma) > 0; \delta_j > 0 (j = 1, \dots, k);$

$$\max \left\{ \left| a_1 z^{\delta_1} \right|, \dots, \left| a_k z^{\delta_k} \right| \right\} < 1 \\ D_z^\nu \{ f(z) \} = \begin{cases} \frac{1}{\Gamma(-\nu)} \int_0^z (z-\xi)^{\nu-1} f(\xi) d\xi & [R(\nu) < 0] \\ \frac{d^n}{dz^n} D_z^{\nu-n} \{ f(z) \} & [0 \leq R(\nu) < n, n \in N_0] \end{cases} \quad (1.8)$$

Generalized elliptic integral of third kind [9]: The generalized form of elliptic integrals is defined by

$$R(\theta, h, \xi; \alpha, t) \\ = \int_0^\theta \frac{1}{(1 + \xi \sin^2 \phi)^\alpha (1 - h^2 \sin^2 \phi)^{1/2-t}} d\phi \quad (1.9)$$

$$R(\theta, h, \xi; \alpha, t) \\ = \int_0^{\sin \theta} \frac{1}{(1 + \xi e^2)^\alpha \sqrt{(1 - e^2)} (1 - h^2 e^2)^{1/2-t}} de, \\ \left(\left| h^2 \right| < 1; 0 \leq \theta \leq \frac{\pi}{2}; t \in C, \alpha \geq 0 \right) \quad (1.10)$$

where ξ is elliptic characteristic and $\xi > -1$. Also, by putting $\alpha = 1$ in (1.9) and (1.10), we have

$$I(\theta, h, \xi; t) = \int_0^\theta \frac{1}{(1 + \xi \sin^2 \phi)(1 - h^2 \sin^2 \phi)^{1/2-t}} d\phi \\ \left(\left| h^2 \right| < 1; 0 \leq \theta \leq \frac{\pi}{2}; t \geq 0 \right) \quad (1.11)$$

$$I(\theta, h, \xi; t) = \int_0^{\sin \theta} \frac{1}{(1 + \xi e^2) \sqrt{(1 - e^2)} (1 - h^2 e^2)^{1/2-t}} de, \\ \left(\left| h^2 \right| < 1; 0 \leq \theta \leq \frac{\pi}{2}; t \geq 0 \right) \quad (1.12)$$

Similarly, on putting $t = 0$ and $\alpha = 1$ in equation (1.9) we get

$$\Pi(\theta, h, \xi) = \int_0^\theta \frac{1}{(1 + \xi \sin^2 \phi)(1 - h^2 \sin^2 \phi)^{1/2}} d\phi, \\ \left(\left| h^2 \right| < 1; 0 \leq \theta \leq \frac{\pi}{2} \right) \quad (1.13)$$

On substituting $\xi = 0$ in equation (1.9), we get elliptic integral of third kind

$$H(\theta, k; \gamma) = \int_0^\theta \frac{1}{(1 - h^2 \sin^2 \phi)^{1/2-t}} d\phi, \\ \left(\left| h^2 \right| < 1; 0 \leq \theta \leq \frac{\pi}{2} \right) \quad (1.14)$$

If we substitute $t = 0, \xi = 0$, then equation (1.9) reduces to $F(\phi, k)$

$$F(\theta, h) = \int_0^\theta \frac{1}{(1 - h^2 \sin^2 \phi)^{1/2}} d\phi, \\ \left(\left| h^2 \right| < 1; 0 \leq \theta \leq \frac{\pi}{2} \right) \quad (1.15)$$

and if $t = 1, \xi = 0$, then our function (1.9) reduces to $E(\theta, h)$

$$E(\theta, h) = \int_0^\theta \sqrt{1 - h^2 \sin^2 \phi} d\phi, \\ \left(\left| h^2 \right| < 1; 0 \leq \theta \leq \frac{\pi}{2} \right). \quad (1.16)$$

Similarly, if we put

$t = 0$ and $\theta = \pi / 2$, then Eq. (1.14) reduces to $K(h) = F(\pi/2, h)$

$$K(h) = \int_0^{\pi/2} \frac{1}{\sqrt{1 - h^2 \sin^2 \phi}} d\phi, \left(\left| h^2 \right| < 1 \right), \quad (1.17)$$

Similarly,

$$E(h) = \int_0^{\pi/2} \sqrt{1-h^2 \sin^2 \phi} d\phi, \quad (|h^2| < 1), \quad (1.18)$$

On applying $k = 3$ in Riemann-Liouville operator defined in (1.7) to the integral in the equation (1.9) with condition $t = \delta - 1 = 1$ and $z = -\sin \theta$, we get

$$R(\theta, h, \xi; \alpha, t) = \sin \theta F_{1;0;0;0}^{1;1;1;1} \left[\begin{matrix} (1:2;2;2)(1/2-t,1); \\ (2:2;2;2); \\ (1/2,1);(\alpha,1) \end{matrix} \middle| h^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \right]. \quad (1.19)$$

Similarly,

$$I(\theta, h, \xi; t) = \sin \theta F_{1;0;0;0}^{1;1;1;1} \left[\begin{matrix} (1:2;2;2)(1/2-t,1); \\ (2:2;2;2); \\ (1/2,1);(1,1) \end{matrix} \middle| h^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \right]. \quad (1.20)$$

With the help of definition of Pochhammer symbol

$$\begin{aligned} \frac{(1)_{2l+2m+2n}}{(2)_{2l+2m+2n}} &= \frac{\Gamma(2l+2m+2n+1)}{\Gamma(2l+2m+2n+2)} \\ &= \frac{\Gamma\left(\frac{1}{2}\right)\left(\frac{1}{2}\right)_{l+m+n}}{2\Gamma\left(\frac{3}{2}\right)\left(\frac{3}{2}\right)_{l+m+n}} = \frac{\left(\frac{1}{2}\right)_{l+m+n}}{\left(\frac{3}{2}\right)_{l+m+n}} \end{aligned} \quad (1.21)$$

where we used the duplication formula defined as follows:

$$\Gamma(2z) = \frac{2^{2z-1}}{\sqrt{\pi}} \Gamma(z) \Gamma\left(z + \frac{1}{2}\right) \quad (1.22)$$

with the help of above relation defined by equation (1.19), we can write as follows:

$$\begin{aligned} R(\theta, h, \xi; \alpha, t) &= \sin \theta F_1 \left[\frac{1}{2}; \frac{1}{2}-t, \frac{1}{2}, \alpha; \frac{3}{2}; h^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \right], \\ &\left(|h^2| < 1; 0 \leq \theta \leq \frac{\pi}{2}; t \in C, \alpha \geq 0 \right). \end{aligned} \quad (1.23)$$

2. Main Results

Theorem1. If Incomplete Fox-Wright Ψ -functions ${}_r\Psi_s^{(\Gamma)}(z)$, is given by

$$\begin{aligned} &{}_r\Psi_s^{(\Gamma)} \left[\begin{matrix} (a_1, \alpha_1, x), (a_j, \alpha_j)_{2,r} \\ (b_j, \beta_j)_{1,s} \end{matrix} \middle| x \right] \\ &= \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \frac{x^n}{n!}. \end{aligned}$$

Under the existence conditions

$$(\alpha_j > 0 \quad (j=1, \dots, r); \beta_j > 0 \quad (j=1, \dots, s);$$

$$1 + \sum_{j=1}^s \beta_j - \sum_{j=1}^r \alpha_j \geq 0).$$

Also let

$$\{\tau, \chi, \omega, \vartheta\} \geq 0, (\tau + \chi > 0; \omega + \vartheta > 0),$$

and

$$\sum_{m=1}^{\infty} \left| \frac{a_m}{m^{1/2(1+\lambda)}} \right| < \infty, \sum_{m=1}^{\infty} \left| \frac{a_m}{m^{1+\mu/2}} \right| < \infty;$$

$$[\tau = 0; R(\lambda) > -1, R(\mu) > -2].$$

Then,

$$\begin{aligned} &\int_0^1 h^\lambda \left(\sqrt{1-h^2} \right)^\mu {}_r\Psi_s^{(\Gamma)} \left[\begin{matrix} (a_1, \alpha_1, y), (a_j, \alpha_j)_{2,r} \\ (b_j, \beta_j)_{1,s} \end{matrix} \middle| xh^\tau \left(\sqrt{1-h^2} \right)^\chi \right] R \left(\theta, \zeta h^\omega \left(\sqrt{1-h^2} \right)^\vartheta, \xi; \alpha, t \right) dh \\ &= \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \\ &\frac{\sin \theta}{2} B \left(\frac{\lambda+1}{2}, \frac{\mu+1}{2} \right) F_{2;m;0;0}^{3;m;1;1;1} \left[\begin{matrix} A: (a_1, \alpha_1) \dots (a_n, \alpha_n); \\ B: (b_1, \beta_1) \dots (b_n, \beta_n); \\ \left(\frac{1}{2}, 1 \right); (\alpha, 1) \end{matrix} \middle| x, \zeta^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \right] \\ &\quad -; -; \end{aligned} \quad (2.1)$$

$$A = \left(\frac{1}{2}; 0, 1, 1, 1 \right), \left(\frac{\lambda+1}{2}, \frac{\tau}{2}, \omega, 0, 0 \right),$$

$$\left(\frac{\mu+2}{2}, \frac{\chi}{2}, \vartheta, 0, 0 \right);$$

$$B = \left(\frac{3}{2}; 0, 1, 1, 1 \right), \left(\frac{\mu+\lambda+3}{2}, \frac{\chi+\tau}{2}, \omega+\vartheta, 0, 0 \right),$$

where $R(\lambda) > -1$ and

$$|\zeta| < 1 \text{ [or } |\zeta| = 1 \text{ and } R(\lambda + 2\omega) > -1].$$

Proof. To prove Theorem 1, we use the series form of Incomplete Fox-Wright Ψ -function given by equation (1.1) and formula of generalized elliptic function of third kind from equation (1.19) in L.H.S. part of theorem, we get

$$\int_0^1 h^\lambda \left(\sqrt{1-h^2} \right)^\mu {}_r\Psi_s^{(\Gamma)} \left[\begin{matrix} (a_1, \alpha_1, y), (a_j, \alpha_j)_{2,r} \\ (b_j, \beta_j)_{1,s} \end{matrix} \right]$$

$$x h^\tau \left(\sqrt{1-h^2} \right)^\chi \left[R \left(\theta, \zeta h^\omega \left(\sqrt{1-h^2} \right)^\vartheta, \xi; \alpha, t \right) dh \right.$$

$$= \int_0^1 h^\lambda \left(\sqrt{1-h^2} \right)^\mu \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \frac{x^n}{n!} h^{\tau n} (1-h^2)^{\frac{n\chi}{2}} \sin \theta$$

$$F_{1;0;0;0}^{1;1;1;1} \left[\begin{matrix} (1; 2, 2, 2) & \left(\frac{1-t}{2}, t, 1 \right) & \left(\frac{1}{2}, 1 \right) & (\alpha, 1) \\ (2; 2, 2, 2) & ; & ; & \end{matrix} \right]$$

$$\zeta^2 h^{2\omega} (1-h^2)^\vartheta \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \Big] dh.$$

Now expanding the multivariable hypergeometric function with the help of (1.5), we get

$$= \int_0^1 h^\lambda \left(\sqrt{1-h^2} \right)^\mu \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1 + \alpha_1 n, y)}{\prod_{j=1}^r \Gamma(b_j + \beta_j n)} \frac{x^n}{n!} h^{\tau n} (1-h^2)^{\frac{n\chi}{2}} \sin \theta$$

$$\frac{(1)_{2n_1+2n_2+2n_3}}{(2)_{2n_1+2n_2+2n_3}} \frac{\left(\frac{1-t}{2} \right)_{n_1} \left(\frac{1}{2} \right)_{n_2} (\alpha)_{n_3}}{n_1! n_2! n_3!} \zeta^{2n_1}$$

$$\zeta^{2n_1} (\sin^2 \theta)^{n_1+n_2+n_3} (-\xi)^{n_3} h^{2\omega n_1} (1-h^2)^{\vartheta n_1} dh.$$

On simplification we get

$$= \frac{1}{2} \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(b_j + \beta_j n)} \frac{x^n}{n!}$$

$$(-\xi)^{n_3} \zeta^{2n_1} (\sin \theta) (\sin^2 \theta)^{n_1+n_2+n_3} \frac{\left(\frac{1}{2} \right)_{n_1+n_2+n_3}}{\left(\frac{3}{2} \right)_{n_1+n_2+n_3}}$$

$$\frac{\left(\frac{1-t}{2} \right)_{n_1} \left(\frac{1}{2} \right)_{n_2} (\alpha)_{n_3}}{n_1! n_2! n_3!} \int_0^1 h^{\lambda+\tau n+2\omega n_1} (1-h^2)^{\frac{\mu}{2} + \frac{n\chi}{2} + \vartheta n_1} dh. \quad (2.2)$$

Now evaluate the integral part using equation (2.2) we arrive

$$= \frac{1}{2} \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(b_j + \beta_j n)} \frac{x^n}{n!}$$

$$(-\xi)^{n_3} \zeta^{2n_1} (\sin \theta) (\sin^2 \theta)^{n_1+n_2+n_3} \frac{\left(\frac{1}{2} \right)_{n_1+n_2+n_3} \left(\frac{1-t}{2} \right)_{n_1} \left(\frac{1}{2} \right)_{n_2} (\alpha)_{n_3}}{\left(\frac{3}{2} \right)_{n_1+n_2+n_3} n_1! n_2! n_3!} \times$$

$$\frac{\Gamma \left(\frac{\lambda+1}{2} \right) \Gamma \left(\frac{\lambda+1}{2} \right)_{\frac{\tau n}{2} + \omega n_1} \Gamma \left(\frac{\mu+1}{2} \right) \Gamma \left(\frac{\mu+1}{2} \right)_{\frac{\chi n}{2} + \vartheta n_1}}{\Gamma \left(\frac{\lambda+\mu+3}{2} \right) \Gamma \left(\frac{\lambda+\mu+3}{2} \right)_{\frac{(\tau+\chi)n}{2} + (\omega+\vartheta)n_1}}.$$

Finally, interpreting the result in terms of multivariable hypergeometric function [10] we obtain the R.H.S. of (2.1).

3. Particular Cases

3.1 Case-1

By using the elliptic function given in equation (1.11), we obtained the following result:

$$\begin{aligned}
 & \int_0^1 h^\lambda \left(\sqrt{1-h^2} \right)^\mu {}_r\psi_s^{(\Gamma)} \left[\begin{matrix} (a_1, \alpha_1, y), (a_j, \alpha_j)_{2,r} \\ (b_j, B_j)_{1,s} \end{matrix} \right] \\
 & x h^\tau \left(\sqrt{1-h^2} \right)^\chi \left[I \left(\theta, \zeta h^\omega \left(\sqrt{1-h^2} \right)^\vartheta, \xi; t \right) dh \right. \\
 & = \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \times \\
 & \frac{\sin \theta}{2} \beta \left(\frac{\lambda+1}{2}, \frac{\mu+1}{2} \right) \times \\
 & F_{2n;0;0;0}^{3n;1;1;1} \left[\begin{matrix} A: (a_1, \alpha_1) \dots (a_n, \alpha_n); \left(\frac{1}{2} - t, 1 \right); \left(\frac{1}{2}, 1 \right); (1, 1) \\ B: (b_1, \beta_1) \dots (b_n, \beta_n); -; -; \end{matrix} \right] \\
 & x, \zeta^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \Big],
 \end{aligned} \tag{3.1}$$

where

$$\begin{aligned}
 A &= \left(\frac{1}{2}; 0, 1, 1, 1 \right), \left(\frac{\lambda+1}{2}, \frac{\tau}{2}, \omega, 0, 0 \right), \\
 & \left(\frac{\mu+2}{2}, \frac{\chi}{2}, \vartheta, 0, 0 \right); \\
 B &= \left(\frac{3}{2}; 0, 1, 1, 1 \right), \left(\frac{\mu+\lambda+3}{2}; \frac{\chi+\tau}{2}, \omega+\vartheta, 0, 0 \right)
 \end{aligned}$$

and $R(\lambda) > -1$ and

$$|\zeta| < 1 \left[\text{or } |\zeta| = 1 \text{ and } R(\lambda + 2\omega) > -1 \right].$$

Proof.

$$\begin{aligned}
 & \int_0^1 h^\lambda \left(\sqrt{1-h^2} \right)^\mu {}_r\psi_s^{(\Gamma)} \left[\begin{matrix} (a_1, \alpha_1, y), (a_j, \alpha_j)_{2,r} \\ (b_j, B_j)_{1,s} \end{matrix} \right] \\
 & x h^\tau \left(\sqrt{1-h^2} \right)^\chi \left[I \left(\theta, \zeta h^\omega \left(\sqrt{1-h^2} \right)^\vartheta, \xi; t \right) dh \right.
 \end{aligned}$$

By using equation (1.1) and (1.20), we get

$$\int_0^1 h^\lambda \left(\sqrt{1-h^2} \right)^\mu \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)}$$

$$\begin{aligned}
 & \frac{x^l}{n!} h^{\tau n} (1-h^2)^{\frac{n\chi}{2}} \sin \theta \times \\
 & F_{1;0;0;0}^{1;1;1;1} \left[\begin{matrix} (1; 2, 2, 2) \quad \left(\frac{1}{2} - t, 1 \right) \left(\frac{1}{2}, 1 \right) \quad (1, 1) \\ (2; 2, 2, 2) \quad ; \quad ; \quad ; \end{matrix} \right] \\
 & \zeta^2 h^{2\omega} (1-h^2)^\vartheta \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \Big] dh.
 \end{aligned}$$

Now using equation (1.5), we have

$$\begin{aligned}
 & = \int_0^1 h^\lambda \left(\sqrt{1-h^2} \right)^\mu \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \times \\
 & \frac{x^l}{n!} h^{\tau n} (1-h^2)^{\frac{n\chi}{2}} \sin \theta \times \frac{(1)_{2n_1+2n_2+2n_3}}{(2)_{2n_1+2n_2+2n_3}} \times \\
 & \frac{\left(\frac{1}{2} - t \right)_{n_1} \left(\frac{1}{2} \right)_{n_2} (1)_{n_3}}{n_1! \, n_2! \, n_3!} \times \\
 & \zeta^{2n_1} (\sin^2 \theta)^{n_1+n_2+n_3} (-\xi)^{n_3} h^{2\omega n_1} (1-h^2)^{9n_1} dh,
 \end{aligned}$$

$$\begin{aligned}
 & = \frac{1}{2} \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \times \\
 & \frac{x^n}{n!} (-\xi)^{n_3} \zeta^{2n_1} (\sin \theta) (\sin^2 \theta)^{n_1+n_2+n_3} \times \\
 & \frac{\left(\frac{1}{2} \right)_{n_1+n_2+n_3}}{\left(\frac{3}{2} \right)_{n_1+n_2+n_3}} \int_0^1 h^{\lambda+\tau n+2\omega n_1} (1-h^2)^{\frac{\mu}{2}+\frac{n\chi}{2}+9n_1} dh.
 \end{aligned}$$

To solve

$$= \int_0^1 h^{\lambda+\tau n+2\omega n_1} (1-h^2)^{\frac{\mu}{2}+\frac{n\chi}{2}+9n_1} dh,$$

$$\begin{aligned}
 \text{let } h^2 = u &\Rightarrow h = u^{1/2} \Rightarrow dh = \frac{1}{2} u^{-1/2} du \\
 &= \int_0^1 u^{\frac{\lambda+\tau n+2\omega n_1}{2}} (1-u)^{\frac{\mu}{2}+\frac{\chi n}{2}+\vartheta n_1} \frac{1}{2} u^{-1/2} du \\
 &= \frac{1}{2} \int_0^1 u^{\frac{\lambda+\tau n+2\omega n_1-1}{2}+1-1} (1-u)^{\frac{\mu}{2}+\frac{\chi n}{2}+\vartheta n_1+1-1} du \\
 &= \frac{1}{2} \frac{\Gamma\left(\frac{\lambda+\tau n+1}{2}+\omega n_1\right) \Gamma\left(\frac{\mu+\chi n+\vartheta n_1}{2}+1\right)}{\Gamma\left\{\left(\frac{\lambda+\tau n+\mu+\chi n+\vartheta n_1+1}{2}\right)+\omega n_1+1\right\}} \\
 &= \frac{1}{2} \frac{\Gamma\left(\frac{\lambda+1}{2}\right) \Gamma\left(\frac{\lambda+1}{2}\right) \frac{\tau n}{2}+\omega n_1 \Gamma\left(\frac{\mu+1}{2}\right) \Gamma\left(\frac{\mu+1}{2}\right) \frac{\chi n}{2}+\vartheta n_1}{\Gamma\left(\frac{\lambda+\mu+3}{2}\right) \Gamma\left(\frac{\lambda+\mu+3}{2}\right) \frac{(\tau+\chi)n}{2}+(\vartheta+\omega)n_1} \\
 &= \frac{1}{2} \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\gamma(a_1+n, y) \prod_{j=2}^r \Gamma(a_j+n)}{\prod_{j=1}^r \Gamma(b_j+l)} \times \\
 &\quad \frac{x^n}{n!} (-\xi)^{n_3} \zeta^{2n_1} (\sin \theta) (\sin^2 \theta)^{n_1+n_2+n_3} \times \\
 &\quad \left(\frac{1}{2}\right)_{n_1+n_2+n_3} \left(\frac{1}{2}-t\right)_{n_1} \left(\frac{1}{2}\right)_{n_2} (1)_{n_3} \times \\
 &\quad \frac{\left(\frac{3}{2}\right)_{n_1+n_2+n_3}}{n_1! \, n_2! \, n_3!} \times \\
 &\quad \frac{\Gamma\left(\frac{\lambda+1}{2}\right) \Gamma\left(\frac{\lambda+1}{2}\right) \frac{\tau n}{2}+\omega n_1 \Gamma\left(\frac{\mu+1}{2}\right) \Gamma\left(\frac{\mu+1}{2}\right) \frac{\chi n}{2}+\vartheta n_1}{\Gamma\left(\frac{\lambda+\mu+3}{2}\right) \Gamma\left(\frac{\lambda+\mu+3}{2}\right) \frac{(\tau+\chi)n}{2}+(\vartheta+\omega)n_1} \\
 &= \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1+\alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j+\alpha_j n)}{\prod_{j=1}^s \Gamma(b_j+\beta_j n)} \times \\
 &\quad \frac{\sin \theta}{2} \beta \left(\frac{\lambda+1}{2}, \frac{\mu+1}{2}\right) \times \\
 &\quad F_{2n;0;0;0}^{3n;1;1;1} \left[\begin{matrix} A: (a_1, \alpha_1) \dots (a_n, \alpha_n); \left(\frac{1}{2}, 1\right); \left(\frac{1}{2}, 1\right); (1, 1) \\ B: (b_1, \beta_1) \dots (b_n, \beta_n); -; -; - \end{matrix} \right] \\
 &\quad x, \zeta^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \Big].
 \end{aligned}$$

3.2 Case-2

By using the elliptic function given in equation (1.13) we obtained the following result:

$$\begin{aligned}
 &\int_0^1 h^\lambda \left(\sqrt{1-h^2}\right)^\mu {}_r \psi_s^{(\Gamma)} \left[\begin{matrix} (a_1, \alpha_1, y), (a_j, \alpha_j)_{2,r} \\ (b_j, B_j)_{1,s} \end{matrix} \right] \\
 &\times x h^\tau \left(\sqrt{1-h^2}\right)^\chi \prod \left(\theta, \zeta h^\omega \left(\sqrt{1-h^2}\right)^\vartheta, \xi; t \right) dh \\
 &= \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1+\alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j+\alpha_j n)}{\prod_{j=1}^s \Gamma(b_j+\beta_j n)} \times \\
 &\quad \frac{\sin \theta}{2} \beta \left(\frac{\lambda+1}{2}, \frac{\mu+1}{2}\right) \times \\
 &\quad F_{2n;0;0;0}^{3n;1;1;1} \left[\begin{matrix} A: (a_1, \alpha_1) \dots (a_n, \alpha_n); \left(\frac{1}{2}, 1\right); \left(\frac{1}{2}, 1\right); (1, 1) \\ B: (b_1, \beta_1) \dots (b_n, \beta_n); -; -; - \end{matrix} \right] \\
 &\quad x, \zeta^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \Big], \\
 &A = \left(\frac{1}{2}; 0, 1, 1, 1\right), \left(\frac{\lambda+1}{2}, \frac{\tau}{2}, \omega, 0, 0\right), \\
 &\left(\frac{\mu+2}{2}, \frac{\chi}{2}, \vartheta, 0, 0\right); \\
 &B = \left(\frac{3}{2}; 0, 1, 1, 1\right), \left(\frac{\mu+\lambda+3}{2}; \frac{\chi+\tau}{2}, \omega+\vartheta, 0, 0\right). \quad (3.2)
 \end{aligned}$$

Proof.

$$\begin{aligned}
 &\int_0^1 h^\lambda \left(\sqrt{1-h^2}\right)^\mu {}_r \psi_s^{(\Gamma)} \left[\begin{matrix} (a_1, \alpha_1, y), (a_j, \alpha_j)_{2,r} \\ (b_j, B_j)_{1,s} \end{matrix} \right] \\
 &\times x h^\tau \left(\sqrt{1-h^2}\right)^\chi \prod \left(\theta, \zeta h^\omega \left(\sqrt{1-h^2}\right)^\vartheta, \xi; t \right) dh.
 \end{aligned}$$

Now by equation (1.1) and (1.13), we get

$$\begin{aligned}
 &= \int_0^1 h^\lambda \left(\sqrt{1-h^2}\right)^\mu \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1+\alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j+\alpha_j n)}{\prod_{j=1}^s \Gamma(b_j+B_j l)} \times \\
 &\quad \frac{x^l}{n!} h^{\tau n} (1-h^2)^{\frac{n\chi}{2}} \sin \theta \times \\
 &\quad F_{1;0;0;0}^{1;1;1;1} \left[\begin{matrix} (1; 2, 2, 2) & \left(\frac{1}{2}, 1\right) & \left(\frac{1}{2}, 1\right) & (1, 1) \\ (2; 2, 2, 2) & ; & ; & ; \end{matrix} \right] \\
 &\quad \zeta^2 h^{2\omega} (1-h^2)^\vartheta \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \Big] dh.
 \end{aligned}$$

Now by equation (1.5), we have

$$\begin{aligned}
 &= \int_0^1 h^\lambda (\sqrt{1-h^2})^\mu \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{l=0}^{\infty} \frac{\Gamma(a_1 + \alpha_l n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^s \Gamma(b_j + \beta_j n)} \times \\
 &\quad \frac{x^n}{n!} h^{\tau n} (1-h^2)^{\frac{n\chi}{2}} \sin \theta \times \frac{(1)_{2n_1+2n_2+2n_3}}{(2)_{2n_1+2n_2+2n_3}} \frac{\left(\frac{1}{2}\right)_{n_1} \left(\frac{1}{2}\right)_{n_2} (1)_{n_3}}{n_1! n_2! n_3!} \times \\
 &\quad \zeta^{2n_1} (\sin^2 \theta)^{n_1+n_2+n_3} (-\xi)^{n_3} h^{2\omega n_1} (1-h^2)^{\mathcal{G}n_1} dh \\
 &= \frac{1}{2} \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1 + \alpha_n n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^s \Gamma(b_j + \beta_j n)} \times \\
 &\quad \frac{z^n}{n!} (-\xi)^{n_3} \zeta^{2n_1} (\sin \theta) (\sin^2 \theta)^{n_1+n_2+n_3} \frac{\left(\frac{1}{2}\right)_{n_1+n_2+n_3}}{\left(\frac{3}{2}\right)_{n_1+n_2+n_3}} \times \\
 &\quad \int_0^1 h^{\lambda+\tau n+2\omega n_1} (1-h^2)^{\frac{\mu}{2}+\frac{n\chi}{2}+\mathcal{G}n_1} dh.
 \end{aligned}$$

To solve

$$= \int_0^1 h^{\lambda+\tau n+2\omega n_1} (1-h^2)^{\frac{\mu}{2}+\frac{n\chi}{2}+\mathcal{G}n_1} dh.$$

$$\text{Let } h^2 = u \Rightarrow h = u^{1/2} \Rightarrow dh = \frac{1}{2} u^{-1/2} du$$

$$\begin{aligned}
 &= \int_0^1 u^{\frac{\lambda+\tau n+2\omega n_1}{2}} (1-u)^{\frac{\mu}{2}+\frac{n\chi}{2}+\mathcal{G}n_1} \frac{1}{2} u^{-1/2} du \\
 &= \frac{1}{2} \int_0^1 u^{\frac{\lambda+\tau n+2\omega n_1-1}{2}+1-1} (1-u)^{\frac{\mu}{2}+\frac{n\chi}{2}+\mathcal{G}n_1+1-1} du \\
 &= \frac{1}{2} \frac{\Gamma\left(\frac{\lambda+\tau n+1}{2}+\omega n_1\right) \Gamma\left(\frac{\mu+\chi n+\mathcal{G}n_1}{2}+1\right)}{\Gamma\left\{\left(\frac{\lambda+\tau n+\mu+\chi n+\mathcal{G}n_1+1}{2}\right)+\omega n_1+1\right\}} \\
 &= \frac{1}{2} \frac{\Gamma\left(\frac{\lambda+1}{2}\right) \Gamma\left(\frac{\lambda+1}{2}\right)^{\frac{\tau n}{2}+\omega n_1} \Gamma\left(\frac{\mu+1}{2}\right) \Gamma\left(\frac{\mu+1}{2}\right)^{\frac{\chi n}{2}+\mathcal{G}n_1}}{\Gamma\left(\frac{\lambda+\mu+3}{2}\right) \Gamma\left(\frac{\lambda+\mu+3}{2}\right)^{\frac{(\tau+\chi)n}{2}+(\mathcal{G}+\omega)n_1}}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{1}{2} \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\gamma(a_1+n, y) \prod_{j=2}^r \Gamma(a_j+n)}{\prod_{j=1}^s \Gamma(b_j+l)} \times \\
 &\quad \frac{x^n}{n!} (-\xi)^{n_3} \zeta^{2n_1} (\sin \theta) (\sin^2 \theta)^{n_1+n_2+n_3} \times \\
 &\quad \frac{\left(\frac{1}{2}\right)_{n_1+n_2+n_3}}{\left(\frac{3}{2}\right)_{n_1+n_2+n_3}} \frac{\left(\frac{1}{2}\right)_{n_1} \left(\frac{1}{2}\right)_{n_2} (1)_{n_3}}{n_1! n_2! n_3!} \times \\
 &\quad \frac{\Gamma\left(\frac{\lambda+1}{2}\right) \Gamma\left(\frac{\lambda+1}{2}\right)^{\frac{\tau n}{2}+\omega n_1} \Gamma\left(\frac{\mu+1}{2}\right) \Gamma\left(\frac{\mu+1}{2}\right)^{\frac{\chi n}{2}+\mathcal{G}n_1}}{\Gamma\left(\frac{\lambda+\mu+3}{2}\right) \Gamma\left(\frac{\lambda+\mu+3}{2}\right)^{\frac{(\tau+\chi)n}{2}+(\mathcal{G}+\omega)n_1}}
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\prod_{j=1}^s \Gamma(b_j)}{\prod_{j=1}^r \Gamma(a_j)} \sum_{n=0}^{\infty} \frac{\Gamma(a_1 + \alpha_n n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^s \Gamma(b_j + \beta_j n)} \times \\
 &\quad \frac{\sin \theta}{2} \beta \left(\frac{\lambda+1}{2}, \frac{\mu+1}{2} \right) \times \\
 &\quad \left| {}_2F_{2,n;0;0}^{3;n;1;1;1} \left[\begin{matrix} A: (a_1, \alpha_1) \dots (a_n, \alpha_n); \left(\frac{1}{2}, 1\right); \left(\frac{1}{2}, 1\right); (1, 1) \\ B: (b_1, \beta_1) \dots (b_n, \beta_n); -; -; \end{matrix} \right] \right| \\
 &\quad x, \zeta^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta]
 \end{aligned}$$

3.3 Case-3

On applying relation given by Srivastava et al. [11][Eq. 6.3] in Theorem 1 we get:

$$\begin{aligned}
 &\int_0^1 h^\lambda (\sqrt{1-h^2})^\mu \gamma_{r,s+1}^{1,r} \left[\begin{matrix} (1-a_1, \alpha_1, y), (1-a_j, \alpha_j)_{2,r} \\ (0, 1), (b_j, \beta_j)_{1,s} \end{matrix} \right] \\
 &\quad - x h^\tau (\sqrt{1-h^2})^\chi \left] R \left(\theta, \zeta h^\omega (\sqrt{1-h^2})^\mathcal{G}, \xi; \alpha, t \right) dh
 \end{aligned}$$

$$\begin{aligned}
 &= \frac{\prod_{j=1}^s \Gamma(b_j) \sum_{n=0}^{\infty} \Gamma(a_1 + \alpha_1 n, y) \prod_{j=2}^r \Gamma(a_j + \alpha_j n)}{\prod_{j=1}^r \Gamma(a_j) \prod_{j=1}^s \Gamma(b_j + \beta_j n)} \times \\
 &\frac{\sin \theta}{2} B\left(\frac{\lambda+1}{2}, \frac{\mu+1}{2}\right) \times \\
 &F_{2n;0;0}^{3n;1;1} \left[\begin{matrix} A: (a_1, \alpha_1) \dots (a_n, \alpha_n); \left(\frac{1}{2}-t, 1\right); \left(\frac{1}{2}, 1\right); (\alpha, 1) \\ B: (b_1, \beta_1) \dots (b_n, \beta_n); -; -; \end{matrix} \right] \\
 &x, \zeta^2 \sin^2 \theta, \sin^2 \theta, -\xi \sin^2 \theta \Big] \\
 &A = \left(\frac{1}{2}; 0, 1, 1, 1\right), \left(\frac{\lambda+1}{2}, \frac{\tau}{2}, \omega, 0, 0\right), \\
 &\left(\frac{\mu+2}{2}, \frac{\chi}{2}, \vartheta, 0, 0\right); \\
 &B = \left(\frac{3}{2}; 0, 1, 1, 1\right), \left(\frac{\mu+\lambda+3}{2}; \frac{\chi+\tau}{2}, \omega+\vartheta, 0, 0\right), \quad (28)
 \end{aligned}$$

where $R(\lambda) > -1$ and

$$|\zeta| < 1 \text{ [or } |\zeta| = 1 \text{ and } R(\lambda + 2\omega) > -1].$$

4. Conclusion and Future Scope

In this work some elliptic integrals of incomplete hypergeometric functions have been established, which are inspired by various applications of different elliptic integrals of complete functions, studied earlier. We have also derived some other relations using the main result in respect of modulus as well as amplitude of the incomplete elliptic integrals. These elliptic integrals can be further used in finding several new integrals pertaining various general functions. These relations can also be employed in derivatives of other properties like analytic continuation and asymptotic behavior.

Reference

- [1] Bansal, M. K., Kumar, D., Khan, I., Singh, J., & Nisar, K. S. (2019). Certain unified integrals associated with product of M-series and incomplete H-functions. *Mathematics*, 7(12), 1191.
- [2] Bansal, M. K., & Kumar, D. (2021). On a family of the incomplete H-functions and associated integral transforms. *Journal of Applied Analysis*, 27(1), 143-52.
- [3] Bansal, M. K., Lal, S., Kumar, D., Kumar, S., & Singh, J. (2020). Fractional differential equation pertaining to an integral operator involving incomplete H-function in the kernel. *Mathematical Methods in the Applied Sciences*.
- [4] Bansal, M. K., Kumar, D., Singh, J., & Nisar, K. S. (2020). Finite and Infinite Integral Formulas Involving the Family of Incomplete H-Functions. *Applications and Applied Mathematics: An International Journal (AAM)*, 15(3), 2.
- [5] Háji, N., Marichev, O. I., & Srivastava, H. M. (1992). A note on the convergence of certain families of multiple hypergeometric series. *Journal of mathematical analysis and applications*, 164(1), 104-15.
- [6] Srivastava, H. M., & Daoust, M. C. (1969). Certain generalized Neumann expansions associated with the Kamp de Friet function, *Nederl. Akad. WetenschIndag. Math*, 31, 449-57.
- [7] Srivastava, H. M., & Daoust, M. C. (1972). A note on the convergence of Kampé de Fériet's double hypergeometric series. *Mathematische Nachrichten*, 53(1-6), 151-9.
- [8] Lin, S. D., Chang, L. F., & Srivastava, H. M. (2009). A certain class of incomplete elliptic integrals and associated definite integrals. *Applied mathematics and computation*, 215(3), 1176-84.
- [9] Dubey, R. S., & Chaurasia, V. B. L. (2013). Definite integrals of generalized certain class of incomplete elliptic integrals. *Tamkang Journal of Mathematics*, 44(2), 197-208.
- [10] Virchenko, N., Kalla, S. L., & Al-Zamel, A. M. A. M. (2001). Some results on a

- generalized hypergeometric function. *Integral Transforms and Special Functions*, 12(1), 89-100.
- [11] Srivastava, H. M. (1995). Some elliptic intervals of Barton and Bushell. *Journal of Physics A: Mathematical and General*, 28(8), 2305.
- [12] Lin, S. D., Tu, S. T., Srivastava, H. M., & Wang, P. Y. (2006). Some families of multiple infinite sums and associated fractional differintegral formulas for power and composite functions. *J. Fract. Calc*, 30, 45-58.
- [13] Srivastava, H. M., Saxena, R. K., & Parmar, R. K. (2018). Some Families of the Incomplete $\overline{H}$ -Functions and Associated Integral Transforms and Operators of Fractional Calculus with Applications. *Russian Journal of Mathematical Physics*, 25(1), 116-38.
- [14] Srivastava, H. M., Chaudhry, M. A., & Agarwal, R. P. (2012). The incomplete Pochhammer symbols and their applications to hypergeometric and related functions. *Integral Transforms and Special Functions*, 23(9), 659-83.
- [15] Srivastava, H. M., Parmar, R. K., & Chopra, P. (2017). Some families of generalized complete and incomplete elliptic-type integrals.
- [16] Dubey, R. S., Sharma, A., & Jain, M. (2018). Study of Incomplete Elliptic Integrals Pertaining to (p) psi (q) Function. *JOURNAL OF APPLIED MATHEMATICS STATISTICS AND INFORMATICS*, 14(2), 11-8.
- [17] Srivastava, H. M., & Bromberg, S. (1995). Some families of generalized elliptic-type integrals. *Mathematical and computer modelling*, 21(3), 29-38.
- [18] Lin, S. D., Chang, L. F., & Srivastava, H. M. (2009). A certain class of incomplete elliptic integrals and associated definite integrals. *Applied mathematics and computation*, 215(3), 1176-84.