

# Antioxidant Activity, Anti-tyrosinase Activity and Fourier Transform Infrared Spectra of Extract and Lotion from Germinated Seeds of *Vigna mungo* (L.) Hepper

Sirikul Thummajitsakul<sup>1,\*</sup>, Phattiya Kriangchaikijkul<sup>1</sup>, Supawee Hrungham<sup>1</sup>, Kittiraphorn Onchai<sup>1</sup>, Kun Silprasit<sup>2</sup>

<sup>1</sup>Faculty of Physical Therapy, Srinakharinwirot University, Nakhon-Nayok 26120, Thailand

<sup>2</sup>Faculty of Environmental Culture and Ecotourism, Srinakharinwirot University, Bangkok 10110, Thailand

Received 19 March 2021; Received in revised form 5 May 2022;

Accepted 18 May 2022; Available online 31 December 2022

## ABSTRACT

This study demonstrated different fingerprints of ethanol extracts, aqueous extracts, and lotion samples from germinated seeds of *Vigna mungo* (L.) Hepper by Fourier transform infrared spectra (FTIR). Moreover, their total phenolic content, antioxidant activity, and tyrosinase inhibition were also determined. The results showed that ethanol and aqueous extracts of germinated *V. mungo* seeds had phenolic content, antioxidant activity, and tyrosinase inhibition. The ethanol extracts had the highest total phenolic content ( $40.77 \pm 7.80$  mg gallic acid equivalents/g extract) and the strongest antioxidant activity ( $1/EC_{50} = 0.009$ ). The high tyrosinase inhibition was found in the lotion base ( $1/EC_{50} = 3.5343$ ), the lotion containing ethanol extracts ( $1/EC_{50} = 1.4473$ ), and aqueous extracts ( $1/EC_{50} = 1.1896$ ). A specific FTIR peak for ethanol extract and lotion samples was assigned to a range of  $2853.42$ - $2979.02$   $\text{cm}^{-1}$ , and two specific peaks for ethanol extracts were at  $1741.65$ - $1742.7$  and  $1539.7$ - $1543.25$   $\text{cm}^{-1}$ . The FTIR peaks of the ethanol extract were observed in 11 wavenumber ranges higher than other samples. Interestingly, the principal component analysis and paired group clustering from the FTIR data, total phenolic content, and biological activities, were able to detect differentiation of the extracts and their lotion samples.

**Keywords:** Antioxidant activity; FTIR; Tyrosinase inhibition; Lotion; *Vigna mungo*

## 1. Introduction

*Vigna mungo* (L.) Hepper belongs to the family Fabaceae, locally known as Black gram, and it is commonly grown in tropical Asian countries [1]. It is one of a number of native legume species in Thailand, which is conserved in the Plant Genetics Conservation Project under the Royal Initiative of Her Royal Highness Princess Maha Chakri Sirindhorn [2]. Its seed is a popular food ingredient as either main course or dessert, and can be consumed as sprouts. Moreover, several studies have reported therapeutic properties of this seed, such as anti-inflammatory, antimicrobial, antioxidant, and antidiabetic activity [3]. Furthermore, it has high nutritional value (i.e., carbohydrates, protein, vitamins, minerals, fat, and fiber) and levels of bioactive compounds (i.e., phenolic compounds, flavonoids, saponins, tannins, alkaloids, and steroids) [4]. Meanwhile, the germination process can further increase its nutritional value and the biochemical activities of its bioactive agents [5-7].

Nowadays, pigmentary disorder is a dermatological condition involving an increased amount of melanin pigments, leading to skin darkening [8]. Tyrosinase is a key enzyme involved in the synthesis of melanins within human melanocytes [9, 10]. Thus, the inhibition of tyrosinase can help to reduce overproduction of melanin pigments, which is useful for cosmetic skin whitening products [11]. It has been reported that some bioactive agents (i.e., phenolics and antioxidants) in plant extracts (i.e., mushroom and riceberry rice) are useful for controlling hyperpigmentation [12-14].

However, several studies of legume seeds focus mainly on nutritional value and bioactive compounds, and development their food products [4, 15-17]. Nowadays, knowledge of black gram sprouts is still limited in cosmetic products of skin whitening. Therefore, our major objectives were to determine the effect of extract and lotion made from germinated *Vigna mungo*

(L.) Hepper seeds on fourier transform infrared spectra (FTIR) fingerprint, total phenolic content, antioxidant activity, and anti-tyrosinase activity.

## 2. Materials and Methods

### 2.1 Chemicals

Chemicals used include: Tyrosinase from mushroom (Sigma), 3,4-Dihydroxy-L-phenylalanine (Sigma), 3-ethylbenzothiazoline-6-sulfonic acid diammonium salt (ABTS) (Sigma), Folin-Ciocalteu's phenol reagent (Merck), Gallic acid (Sigma-Aldrich), Absolute ethanol (Sigma-Aldrich), Potassium persulfate (Ajax Finechem), 1-Hexadecanol (Sigma-Aldrich), Potassium sorbate (Sigma-Aldrich), Emulsifying wax (Sigma-Aldrich), and Glycerol (Sigma-Aldrich).

### 2.2 Preparation of germinated seeds

*Vigna mungo* (L.) Hepper was purchased from a market and its external morphology was identified by comparing it with the samples from the E-learning of the Department of Agronomy, Kasetsart University, Kamphaengsean Campus [40]. For the germination process, the black gram was cleaned with water and then soaked in water for 24 hours. Following this, the outer layer of the germinated seeds were peeled off. After peeling, the samples were incubated at 50°C for 16 hours or until dry. Then, the dried samples were ground into fine powder and kept at 4°C until use.

### 2.3 Extraction of ground samples

Ground sample (10 g) was extracted with 250 ml solvent (absolute ethanol and distilled water) for 24 hours at 45 °C. The extraction process was performed twice. Each extract was sieved *via* filter cloth. Then, each filtered extract was evaporated at 45°C for either 19 mins (absolute ethanol extract) or 40 mins (aqueous extract) using a rotary evaporator (IKAa RV10) according to the protocol of Thummajitsakul et al. (2019) [18]. A gummy extract was obtained and

further diluted with its respective solvent for final concentration of 10 mg/ml.

## 2.4 Lotion preparation

A lotion base was prepared by mixing 3% cetyl alcohol, 3% emulsifying wax, 4% coconut oil, and 2% polysorbate 80 at 70 °C until the mixture was clear. Then, 2% glycerol was added to the mixture and stirred until its temperature decreased to 40 °C. For the lotion formulation, each sample of germinated *V. mungo* seed extract (10 ml) was added to the lotion base and left to cool down to room temperature.

## 2.5 Total phenolic contents

The total phenolic content of each extract was estimated by using the Folin-Ciocalteu method [18]. Each extract (300 µl) was thoroughly mixed with 1.5 ml of Folin-Ciocalteu reagent for 5 minutes at room temperature, followed by mixing with sodium carbonate (7.5% w/v) for a final volume of 1.2 ml, and allowed stand for 30 minutes at room temperature. The absorbance of each reaction was detected at 765 nm. Each reaction was carried out 2 times. Gallic acid (0 - 1 mg/ml) was used to generate the standard curve ( $R^2=1.00$ ). The total phenolic content of each extract was expressed as mg of gallic acid equivalent per g extract.

## 2.6 3-ethylbenzothiazoline-6-sulfonic acid diammonium salt (ABTS)

Antioxidant activity of each sample was evaluated using the ABTS free-radical scavenging method [18]. The ABTS radical cation solution was carried out by the reaction of 7mM ABTS (10 ml) and 140 mM potassium persulfate (179 µl) under darkness at room temperature for 16 hours. Then, the ABTS radical cation solution was diluted with distilled water until its absorbance at 734 nm measured  $0.700 \pm 0.050$ . Each extract (20 µl) was then reacted in darkness with the diluted ABTS radical cation solution (3.9 ml). The absorbance of

each reaction was determined after 6 minutes at 734 nm. Each reaction was performed in duplicate. The percentage of antioxidant activity was measured using the equation below:

$$\% \text{Antioxidant capacity} = \frac{(OD_{ABTS} - OD_{\text{Sample-ABTS}})}{OD_{ABTS}} \times 100\% \quad (2.1)$$

$OD_{ABTS}$  is the absorbance of the diluted  $ABTS^{*+}$  solution, while  $OD_{\text{Sample-ABTS}}$  is the absorbance of a mixture of the diluted  $ABTS^{*+}$  solution and each sample.

The percentage of antioxidant activity of each sample was used to generate a simple linear regression ( $R^2=0.95-1.00$ ) to determine the effective concentration to scavenge 50% of radicals ( $EC_{50}$ ).

## 2.7 Tyrosinase inhibition assay

Tyrosinase inhibitory activity was determined using 3-(3,4-dihydroxyphenyl)-L-alanine (L-DOPA) as a substrate [19]. Briefly, 5 mM L-DOPA (100 µl), 0.1M sodium phosphate buffer pH 6.8 (20 µl), and seed extract (40 µl) were mixed. For the negative control, sodium phosphate buffer was used in place of extract. Then to each mixture, 40 µl of mushroom tyrosinase (200 Units/ml) was added, and then incubated at 37°C for 20 min. The absorption was determined at 450 nm. Kojic acid (10 mg/ml) was used as positive control. The percentage of tyrosinase inhibition was calculated using the equation below:

$$\% \text{Tyrosinase inhibition} = \frac{(OD_{\text{blank}} - OD_{\text{sample}})}{OD_{\text{blank}}} \times 100, \quad (2.2)$$

where  $OD_{\text{blank}}$  is the absorbance of each reaction without each sample at 450 nm, and  $OD_{\text{sample}}$  is the absorbance of each reaction with each sample at 450 nm.

The percentage of tyrosinase inhibition of each sample was used to

generate a simple linear regression ( $R^2=0.70-1.00$ ) to determine the effective concentration to inhibit 50% of tyrosinase ( $EC_{50}$ ).

## 2.8 Fourier transform infrared spectra (FTIR) fingerprint

FTIR peaks in aqueous extracts, ethanol extracts, and lotion samples were detected in a range of 550 to 4000  $\text{cm}^{-1}$  and a resolution of 4  $\text{cm}^{-1}$  by FTIR spectroscopy (PerkinElmer spectrum IR version 10.6.0). Each sample was measured in duplicate [20-27].

## 2.9 Statistical analysis

PSPP version 0.10.5 [28] and the paleontological statistic program version 3.16 [29] were used for statistical analysis. Descriptive statistical analysis of total phenolic contents, antioxidant activity, and tyrosinase inhibitory activity is expressed as mean, standard deviation, and percentage. Moreover, the difference in total phenolic content, antioxidant activity, tyrosinase inhibitory activity, and FTIR data among sample groups were all analyzed using one-way analysis of variance (one-way ANOVA). Principal component analysis (PCA) was used to summarize the data, while paired group (UPGMA) clustering (bootstrap=10000) was applied to express differentiation of the samples.


## 3. Results and Discussion

The functional plant ingredients and product quality can be directly or indirectly influenced by the production process. It has been reported that legumes consist of various phytochemicals that are beneficial

for human health [4, 17]. The germination process especially effects the level of bioactive compounds and antioxidants in seeds, with a germination period of 1-5 days being best for increasing the level of these compounds [30].

## 3.1 Fourier transform infrared spectra (FTIR) fingerprint

In current study, the FTIR fingerprints of ethanol extracts, aqueous extracts, powder sample, and lotion samples of germinated *V. mungo* seeds showed differences in the range of 550 to 4000  $\text{cm}^{-1}$  (Fig. 1). The results demonstrated the effect of extraction solvent and lotion samples of *V. mungo* on the FTIR fingerprints. The FTIR peaks in 12 wavenumber ranges for each sample were identified and shown in Table 1. The results showed a specific peak for ethanol extract and lotion samples in the range of 2853.42-2979.02  $\text{cm}^{-1}$ , which indicate C–H stretching in  $\text{CH}_2$  groups (carbohydrates and fats) [20, 25, 26], while specific peaks for ethanol extract were at 1741.65-1742.7 and 1539.7-1543.25  $\text{cm}^{-1}$ , which indicate C=O in lipids and N–H of proteins, respectively [22, 24, 26]. Interestingly, the greatest number of peaks were found in the ethanol extract for 11 wavenumber ranges, followed by aqueous extracts (8 wavenumber ranges), lotion samples (5 and 8 wavenumber ranges), and powder sample (4 wavenumber ranges) of germinated *V. mungo* seeds.



**Fig. 1.** Example of FTIR spectra of extracts, lotion samples, and powder sample of germinated *Vigna mungo* seeds. A: ethanol extract, B: aqueous extract, C: powder sample, D: lotion sample with ethanol extract, E: lotion sample with aqueous extract. Peak a is specific band for ethanol extract, ethanol solvent, and lotion samples, while peaks b and c are specific bands for ethanol extract.

**Table 1.** Wavenumber ranges of FTIR peaks and functional groups of ethanol extracts, aqueous extracts, and powder sample of germinated *V. mungo* seeds.

Peak number	Wavenumber range of extract and powder samples in this study (cm <sup>-1</sup> )	Wavenumber range of references (cm <sup>-1</sup> )	Assignment	Function groups	References
1	3011.58-3339.80	3000–3600	O-H stretch	water, alcohols, phenols, carbohydrates, peroxides	[22, 25]
2	2853.42 -2979.02	3000–2800	C–H stretching of CH <sub>2</sub> groups	carbohydrates and fats	[20, 25, 26] [25, 26]
3	1632.44-1742.7	1600-1760	N-H bending vibrations, C=O bending vibrations	amino acids, fatty acids, ester	[26]
4	1741.65-1742.7	1700-1799	C=O of lipids	lipids	[24]
5	1632.44-1647.3	1600-1706	Amide I of proteins, C=O	proteins	[24]
6	1380.81-1456.05	1380-1465	CH <sub>3</sub> lipids/proteins and COO– of amino acids	lipids/proteins/ amino acids	[24]
7	1539.7-1543.25	1460-1590	Amide II of proteins, N–H	proteins	[24]
8	1328.24-1411.19	1300-1450	Primary or secondary O–H bending (in-plane), and phenol or tertiary alcohol (O–H bend)	phenyl groups	[21, 22]
9	1328.24-1377.5	1300-1380	CH <sub>3</sub> bending	lipid	[23]
10	1150.3-1229.92	1150-1270	C–O stretching vibrations	acid or ester	[22, 26]
11	1016.43-1047.52	1015-1080	glycosidic	carbohydrates	[27]
12	507.24-998.71	< 1000	C-H bending vibrations	Isoprenoids	[22, 26]

### 3.2 Total phenolic content

Total phenolic content, antioxidant activity, and tyrosinase inhibitory activity were determined in ethanol extracts, aqueous extracts, lotion containing either aqueous or ethanol extract, and finally lotion base. The results showed that the highest level ( $p < 0.05$ ) of total phenolic content was significantly observed in ethanol extracts ( $40.77 \pm 7.80$  mg gallic acid equivalents / g extract), followed by aqueous extract ( $17.91 \pm 2.73$  mg gallic acid equivalents / g extract). For the lotion samples, the highest level of total phenolic content was found in lotion containing ethanol extract ( $2.18 \pm 0.74$  mg gallic acid equivalents / mL lotion), followed by lotion containing aqueous extract ( $0.86 \pm 0.20$  mg gallic acid equivalents / mL lotion), and lotion base ( $0.75 \pm 0.17$  mg

gallic acid equivalents / mL lotion) ( $p$ -value $<0.05$ ) (Table 2).

### 3.3 Antioxidant activity and tyrosinase inhibitory activity

Antioxidant activity of ethanol extract and aqueous extract showed low  $1/EC_{50}$  values ( $1/EC_{50} = 0.009$  and  $0.005$ , respectively). Similarly, ethanol extract and aqueous extract showed tyrosinase inhibitory activity with low  $1/EC_{50}$  values ( $1/EC_{50} = 0.0454$  and  $0.0025$ , respectively). However, antioxidant activity was not observed in all lotion samples; tyrosinase inhibitory activity was found at the highest level in lotion base ( $1/EC_{50} = 3.5343$ ), followed by lotion containing ethanol extract ( $1/EC_{50} = 1.4473$ ), and then lotion containing aqueous extract ( $1/EC_{50} = 1.1896$ ) ( $p$ -value $<0.05$ ) (Table 2).

**Table 2.** Total phenolic contents, antioxidant activities, tyrosinase inhibition, and the number of FTIR peaks of ethanol extracts, aqueous extracts, lotion samples and powder samples of germinated *V. mungo* seeds.

Samples	Methods of sample preparation	Total phenolic contents (mg gallic acid equivalents / g extract) **	Antioxidant activities		Tyrosinase inhibition		The number of FTIR peaks
			EC <sub>50</sub> (mg/ml)	1/EC <sub>50</sub>	EC <sub>50</sub> (mg/ml)	1/EC <sub>50</sub>	
Ethanol	extract	40.77±7.80	105.27±30.94	0.009	22.02±6.74	0.0454	11
	lotion	2.18±0.74	0	-	0.69±0.07	1.4473	8
Aqueous	extract	17.91±2.73	195.98±92.06	0.005	398.86 ±243.29	0.0025	8
	lotion	0.86±0.20	0	-	0.84±0.19	1.1896	5
Lotion base	lotion	0.75±0.17	0	-	0.28±0.05	3.5343	5
Powder samples	grinding	-	-	-	-	-	4
Kojic acid (25-100 mg/ml)					132.73±14.34	0.0076	
P-value		0.000*	0.000*		0.001*		

**Note:** \*Differentiation of total phenolic content, antioxidant activity, and tyrosinase inhibition among ethanol extract and aqueous extract, and lotion samples of germinated *V. mungo* seeds were measured at a significance level of 0.05 (*P*-value <0.05).

\*\* Total phenolic content of lotion samples are expressed as mg gallic/ml lotion

### 3.4 Principal component analysis (PCA) and paired group (UPGMA) clustering

PCA analysis was carried out on FTIR data in the 550 to 4000 cm<sup>-1</sup> regions (Figure 2A). The results are shown as scatter plots of the first and second principal components, wherein PC1 contained 45.40% and PC2 contained 24.77% of the spectra variability. The PCA biplot was able to distinguish the FTIR spectrum among samples of the ethanol extracts, aqueous extracts, powder sample, and lotion samples. Additionally, the PCA results indicated that the powder sample, lotion base, and lotion containing aqueous extract had similar chemical compositions. Moreover, aqueous extract and lotion containing ethanol extract had similarity to each other, while the ethanol extract was distinguished from the other samples on the FTIR spectrum (Fig. 2A). Corresponding to the results of PCA analysis, cluster analysis confirmed differentiation of the ethanol extract from the other samples (Figs. 2A and 3A). The ethanol extracts were distributed in positive loadings plot of PC1. This plot demonstrates the major peaks found in the ethanol extract at approximately 1150.3-1229.92, 1328.24-1377.5, 1539.7-1543.25, 1741.65-1742.7, and 2918.55-2979.02 cm<sup>-1</sup>. Of these bands, specific peaks for ethanol

extracts were seen at 1741.65-1742.7 and 1539.7-1543.25 cm<sup>-1</sup>.


Additionally, the aqueous extract and lotion containing ethanol extract were distributed in negative loadings plot of PC1. This plot demonstrates significant peaks found in the ethanol extract at approximately 1380.81-1456.05, 1328.24-1411.19, and 1016.43-1047.52 cm<sup>-1</sup>. For the powder sample, lotion base, and lotion containing aqueous extract, peaks were distributed in negative loadings plot of PC2.

The results confirmed that the PCA analysis was able to detect differentiation of the FTIR spectrum of extractions and their products, which was consistent with the cluster analysis. It indicated that extraction solvent and lotion samples had an effect on FTIR spectra.

As the results in Figure 2B shows, PC1 and PC2 were 87.03% and 12.71% of the data variance of total phenolic content, antioxidant activity, and tyrosinase inhibitory activity. The PCA analysis confirmed that the highest total phenolic content and antioxidant activity was found in the ethanol extract, followed by the aqueous extract, while the highest level of tyrosinase inhibitory activity was found in lotion base, lotion containing ethanol extract, and lotion containing aqueous extract.


Similarly, cluster analysis showed differentiation between the ethanol extract and the other samples (Fig. 3B), corresponding to the cluster analysis of FTIR data (Fig. 3A). However, antioxidant activity showed the strongest positive correlation with total phenolic content ( $r = 0.989$ ,  $p$ -value $<0.05$ ), while the correlation between antioxidant activity and tyrosinase inhibition

was insignificant ( $r = -0.909$ ,  $p$ -value $>0.05$ ), as were the correlations between antioxidant activity and FTIR data ( $r = 0.845$ ,  $p$ -value $>0.05$ ), between tyrosinase inhibition and total phenolic content ( $r = -0.840$ ,  $p$ -value $>0.05$ ), between tyrosinase inhibition and FTIR data ( $r = -0.571$ ,  $p$ -value $>0.05$ ), and between FTIR data and total phenolic content ( $r = 0.899$ ,  $p$ -value $>0.05$ ).



**Fig. 2.** Principal component analysis (PCA). A: PCA biplot from FTIR data, B: PCA biplot from total phenolic contents, antioxidant activities, and tyrosinase inhibition of ethanol extracts, aqueous extracts, and lotion samples of germinated *V. mungo* seeds. LVMW and LVME are lotion from aqueous extract and ethanol extract, respectively. VMW and VME are aqueous extract and ethanol extract, respectively. VMP is powder sample of germinated *V. mungo* seeds.





**Fig. 3.** Paired group (UPGMA) clustering (bootstrap=10000) of extract and powder samples of germinated seeds of *V. mungo*, and each solvent. A was UPGMA clustering from FTIR data, and B was UPGMA clustering from total phenolic contents, antioxidant activities, and tyrosinase inhibition of ethanol extracts, aqueous extracts, and lotion samples of germinated seeds of *V. mungo*.

The FTIR spectrum of germinated seed powder, all extracts, and lotion samples provided interesting results involving the presence of bioactive compounds. Several bioactive compounds provided different functional groups. These functional groups may play very important roles in enhancement of the antioxidant activity and tyrosinase inhibitory activity of the extracts and lotion samples of germinated *V. mungo* seeds. Previously, it has been reported that some bioactive compounds with a higher phenolic hydroxyl number have greater antioxidant and tyrosinase inhibitory activities (i.e., the most activity is in dihydromyricetin, followed by rosmarinic acid, baicalein, shikonin, then isoeugenol) [31]. Phenol hydroxyl compounds (i.e., flavonoids) have a similar structure to tyrosinase substrates (i.e., tyrosine and dopamine), which can affect the inhibition of tyrosinase activity [32]. A greater number of phenolic hydroxyl groups can influence the antioxidant activity by providing more  $H^+$  to scavenge free radicals and hydrogen bonds [33]. Moreover, organic solvent may provide a greater number of phenolic compounds with active functional groups than aqueous solvent does. It has been reported that absolute ethanol extract can provide the greatest antioxidant activity, compared to those of aqueous solvent or ethanol solvents with increasing water content [34].

Previously, a study on FTIR spectroscopy in extracts of *Garcinia schomburgkiana* leaves showed that four specific functional groups are found in ethanol extract, while only one functional group is found in aqueous extract [41].

Moreover, some FTIR peaks seen in the lotion samples matched with those of the ethanol and aqueous extracts, demonstrating some FTIR peaks of extracts and powder sample being revealed in lotion samples. The results showed the presence of phenolic compounds in extracts and lotion samples of germinated *V. mungo* seeds. Phenolic compounds are responsible for various

medicinal properties, such as antioxidant, anti-tyrosinase, and anti-diabetic activity [32, 33, 35]. Additionally, it has been reported that the antioxidant and tyrosinase activities of medicinal plant extracts are related to total phenolic contents [36].

However, the lotion samples showed total phenolic content and antioxidant activity lower than the ethanol and aqueous extracts, while the lotion samples showed more effective tyrosinase inhibitory activity than the extracts did. The efficiency of solvent extraction may affect solubility of multiple agents in lotion and plant extract.

However, in this study, the lotion sample contained different ingredients, namely cetyl-alcohol, emulsifying wax, coconut oil, polysorbate 80, and plant extract. Among the ingredients, coconut oil is commonly applied as an important ingredient in cosmetics such as cream and lotion and has been shown to have several biological activities such as anti-inflammatory and skin protective properties [37].

Virgin coconut oil contains several fatty acids such as lauric acid, myristic acid, palmitic acid, and linoleic acid. Among the fatty acids, linoleic acid shows a skin whitening effect [38]. The combination of various active components in the lotion base can provide greater activity than the sum of each component. For example, it has been reported that repellent cream formulations, which are prepared from different essential oils, can provide a synergistic benefit against *Culex quinquefasciatus* and *Aedes aegypti* [39]. Moreover, the anti-tyrosinase activity in the lotion base was found to be greater than the lotions containing plant extract. It indicates that the lotion base employed in this current study was not suitable since some ingredients in the lotion base may interfere with tyrosinase inhibitory activity.

#### 4. Conclusion

In conclusion, this study confirmed that ethanol and aqueous extracts of germinated *V. mungo* seeds had phenolic content, antioxidant activity, and tyrosinase inhibitory activity. Moreover, the lotion base, lotion containing aqueous extract, and lotion containing ethanol extract showed tyrosinase inhibitory activity higher than that of the extracts. Among all lotion formulations studied, the best tyrosinase inhibitory activity was found in the lotion base. Moreover, powder germinated seeds, ethanol extracts, aqueous extracts, and lotion samples provided different FTIR fingerprints. The PCA and UPGMA clustering, obtained from the FTIR data, total phenolic content, and biological activities, were able to differentiate between extracts and their lotion samples. The FTIR data of germinated seed powder, all extracts, and lotion samples showed the presence of several bioactive compounds. The bioactive compounds with different functional groups may involve the antioxidant activity and tyrosinase inhibitory activity of the extracts of germinated *V. mungo* seeds. Thus, it can be concluded that these effective seed extracts can be applied for the formulation of skin whitening lotion.

#### Acknowledgements

The authors gratefully acknowledge a grant support (Grant number: 272/2563) by Faculty of Physical Therapy, Srinakharinwirot University, and research tools by Faculty of Environmental Culture and Ecotourism of Srinakharinwirot University. The authors declare that there are no conflicts of interest.

#### References

- [1] Tomooka N, Kaga A, Vaughan DA. 2006. The Asian *Vigna* (*Vigna* subgenus *Ceratotropis*) biodiversity and evolution. In: Sharma AK, Sharma S, editors. Plant genome, biodiversity and evolution. Phanerogams (*Angiosperms* *Dicotyledons*). 1st ed. Science Publishers, NJ:New Jersey; 2006. p. 87-126.
- [2] Chotiyarnwong O, Reanhatthakam K, Chanmuang A, Chareonsap P, Jutamas P. Thailand native legume genetic resources in Plant Genetics Conservation Project under the Royal initiative of Her Royal Highness Princess Maha Chakri Sirindhorn. Kaen Kaset 2011; 39(3): 319-27.
- [3] Zaheer M, Ahmed S, Hassan MM. A review of medicinal uses, phytochemistry and pharmacology of *Vigna mungo* (L.) Hepper. J Pharmacogn Phytochem 2020; 9(1):1307-9.
- [4] Varma RK, Garg VK, Singh L, Kumar D. Pharmacognostic evaluation and phytochemical analysis of seeds of *Vigna mungo* (L.) Hepper. Open Res J Phytother Pharmacognosy 2013; 1(1): 1-9.
- [5] Vidal-Valverde C, Frias J, Sierra I, Blazquez I, Lambein F, Kuo Y. New functional legume foods by germination: Effect on the nutritive value of beans, lentils and peas. Eur Food Res Technol 2002;215:472-7.
- [6] Xu M, Dong J, Zhu M. Effects of germination conditions on ascorbic acid level and yield of soybeans sprouts. J Sci Food Agric 2005;85:943-7.
- [7] Tiansawang K, Luangpituksa P, Varayanond W, Hansawasdi C. GABA ( $\gamma$ -aminobutyric acid) production, antioxidant activity in some germinated dietary seeds and the effect of cooking on their GABA content. Food Sci Technol 2016; 36(2):313-21.
- [8] Halder RM, Nandedkar MA, Neal KW. Pigmentary disorders in ethnic skin. Dermatol Clin 2003;21:617-28.
- [9] Schallreuter KU, Salem MMAEL, Hasse S, Rokos H. The redox-biochemistry of human hair pigmentation. Pigment Cell Melanoma Res 2011; 24(1):51-62.

- [10] Zaidi KU, Ali SA, Ali AS. Effect of purified mushroom tyrosinase on melanin content and melanogenic protein expression. *Biotechnol Res Int* 2016; 2016:1-8.
- [11] Lai J, Xin C, Zhao Y, Feng B, He C, Dong Y, Fang Y, Wei S. Study of active ingredients in black soybean sprouts and their safety in cosmetic use. *Molecules* 2012; 17:11669-79.
- [12] An SM, Koh JS, Boo YC. p-Coumaric acid not only inhibits human tyrosinase activity in vitro but also melanogenesis in cells exposed to UVB. *Phytother Res* 2010;24:1175-80.
- [13] Im KH, Baek SA, Choi J, Lee TS. Antioxidant, anti-melanogenic and anti-wrinkle effects of *Phellinus vaninii*. *Mycobiology* 2019;47:494-505.
- [14] Rodboon T, Okada S, Suwannalert P. Germinated riceberry rice enhanced protocatechuic acid and vanillic acid to suppress melanogenesis through cellular oxidant-related tyrosinase activity in B16 Cells. *Antioxidants (Basel)* 2020; 9(3):247.
- [15] Castrorubio A, Garcia M, Marina M. Rapid separation of soybean and cereal (wheat, corn, and rice) proteins in complex mixtures: Application to the selective determination of the soybean protein content in commercial cereal-based products. *Anal Chim Acta* 2006; 558(1-2):28-34.
- [16] Girish TK, Pratapa VM, Prasada Rao UJS. Nutrient distribution, phenolic acid composition, antioxidant and alphaglucosidase inhibitory potentials of black gram (*Vigna mungo* L.) and its milled by-products. *Food Res Int* 2012; 46(1):370-7.
- [17] Zaheer M, Ahmed S, Hassan MM. A review of medicinal uses, phytochemistry and pharmacology of *Vigna mungo* (L.) Hepper. *J Pharmacogn Phytochem* 2020; 9(1):1307-9.
- [18] Thummajitsakul S, Boonburapong B, Silprasit K. Antioxidant and antidiabetic effects of *Garcinia schomburgkiana* extracts and fermented juices. *Pertanika J Trop Agric Sci* 2019;42(1):45-60.
- [19] Liang C, Lim JH, Kim SH, Kim DS. Dioscin: A synergistic tyrosinase inhibitor from the roots of *Smilax china*. *Food Chem* 2012; 134(2):1146-8.
- [20] Manrique GD, Lajolo FM. FTIR spectroscopy as a tool for measuring degree of methyl esterification in pectins isolated from ripening papaya fruit. *Postharvest Biol Technol* 2002;25:99-107.
- [21] Coates J. Interpretation of infrared spectra, a practical approach. In: Meyers RA, editors. *Encyclopedia of analytical chemistry*. John Wiley & Sons Ltd. , UK:Chichester, 2006. p. 10815-37.
- [22] Caunii A, Pribac G, Grozea I, Gaitin D, Samfira I. Design of optimal solvent for extraction of bioactive ingredients from six varieties of *Medicago sativa*. *Chem Cent J* 2012;6:123.
- [23] Baker MJ, Trevisan J, Bassan P, Bhargava R, Butler HJ, Dorling KM, Fielden PR, Fogarty SW, Fullwood NJ, Heys KA, Hughes C, Lasch P, Martin-Hirsch PL, Obinaju B, Sockalingum GD, Sulé-Suso J, Strong RJ, Walsh MJ, Wood BR, Gardner P, Martin FL. Using fourier transform IR spectroscopy to analyze biological materials. *Nat Protoc* 2014;9:1771-91.
- [24] Hands JR, Clemens G, Stables R, Ashton K, Brodbelt A, Davis C, Dawson TP, Jenkinson MD, Lea RW, Walker C, Baker MJ. Brain tumour differentiation: rapid stratified serum diagnostics via attenuated total reflection Fourier-transform infrared spectroscopy. *J Neurooncol* 2016; 127: 463-72.
- [25] Cao Z, Wang Z, Shang Z, Zhao J. Classification and identification of *Rhodobryum roseum* Limpr. and its adulterants based on fourier-transform

- infrared spectroscopy (FTIR) and chemometrics. PLoS One 2017; 12(2): e0172359.
- [26] Topalaa CM, Tatarua LD. ATR-FTIR spectra fingerprinting of medicinal herbs extracts prepared using microwave extraction. Arabian J Med Aroma Plants 2017;3:1-9.
- [27] Duraka T, Depciuch J. Effect of plant sample preparation and measuring methods on ATR-FTIR spectra results. Environ Exp Bot 2020;169:103915.
- [28] Pfaff B, Darrington J, Stover J, Satman MH, Beckmann F, Williams J, Kieffe M, Kobly P, van Son R. 2013. GNU PSPP version [Internet]. [cited 2021 Jan 20]. Available from: <https://www.gnu.org/software/pspp/>
- [29] Hammer Ø, Harper DAT, Ryan PD. PAST: paleontological statistics software package for education and data analysis. Palaeontol Electronica 2001;4(1):1-9.
- [30] Xue Z, Wang C, Zhai L, Yu W, Chang H, Kou X, Zhou F. Bioactive compounds and antioxidant activity of mung bean (*Vigna radiata* L.), soybean (*Glycine max* L.) and black bean (*Phaseolus vulgaris* L.) during the germination process. Czech J Food Sci 2016;34:68-78.
- [31] Zuo AR, Dong HH, Yu YY, Shu QL, Zheng LX, Yu XY, Cao SW. The antityrosinase and antioxidant activities of favonoids dominated by the number and location of phenolic hydroxyl groups. Chin Med 2018;13:51.
- [32] Thomeby-andersson K, Sterner O, Hansson C. Tyrosinase-mediated formation of a reactive quinone from the depigmenting agents, 4-tert-butylphenol and 4-tert-butyl-catechol. Pigment Cell Res. 2000;13:33-8.
- [33] Cai YZ, Sun M, Xing J. Structure-radical scavenging activity relationships of phenolic compounds from traditional Chinese medicinal plants. Life Sci 2006; 78:2872-88.
- [34] Do QD, Angkawijaya AE, Tran-Nguyen PL, Huynh LH, Soetaredjo FE, Ismadji S, Ju YH. Effect of extraction solvent on total phenol content, total flavonoid content, and antioxidant activity of *Limnophila aromatica*. J Food Drug Anal 2014; 22(3): 296-302.
- [35] You Q, Chen F, Wang X, Jiang Y, Lin S. Anti- diabetic activities of phenolic compounds in muscadine against alpha-glucosidase and pancreatic lipase. LWT 2012; 46(1):164-8.
- [36] Muddathir AM, Amiuchi KY, Batubara I, Mohieldin EAM, Mitsunag T. Anti-tyrosinase, total phenolic content and antioxidant activity of selected Sudanese medicinal plants. S Afr J Bot 2017; 109:9-15.
- [37] Varma SR, Sivaprakasam TO, Arumugam I, Dilip N, Raghuraman M, Pavan KB, Rafiq M, Paramesh R. In vitro anti-inflammatory and skin protective properties of Virgin coconut oil. J Tradit Complement Med 2019;9(1):5-14.
- [38] Shigeta Y, Imanaka H, Ando H, Ryu A, Oku N, Baba N, Makino T. Skin whitening effect of linoleic acid is enhanced by liposomal formulations. Biol Pharm Bull 2004; 27(4):591-4
- [39] Reegan AD, Kannan RV, Paulraj MG, Ignacimuthu S. Synergistic effects of essential oil-based cream formulations against *Culex quinquefasciatus* Say and *Aedes aegypti* L. (Diptera: Culicidae). J Asia Pac Entomol 2014;17(3):327-31.
- [40] E-learning. Department of Agronomy Kasetsart University Kamphaengsean Campus [Internet]. [cited 2020 Nov 11]. Available from: <http://agron.agri.kps.ku.ac.th/index.php/th/2015-04-20-02-02-25/30-economic-crops>

- [41] Thummajitsakul S, Samaikam S, Tacha S, Silprasit K. Study on FTIR spectroscopy, total phenolic content, antioxidant activity and anti-amylase activity of extracts and different tea forms of *Garcinia schomburgkiana* leaves. LWT. 2020;134:110005.