



Investigation of Larvicidal Activity and Histopathological Variations of Brown Algae *Spatoglossum asperum* J. Agardh against *Aedes aegypti*, *Culex quinquefasciatus*, and *Anopheles stephensi*

Baluchamy Poornasundari^{1,*}, Subramanian Arivoli¹, Dinakararajan Pradiba²,
Samykanu Mariaamalraj³, Kumaresan Velmurugan⁴

¹Department of Zoology, Thiruvalluvar University, TamilNadu 632115, India

²Department of Biotechnology, Kamaraj College of Engineering & Technology, TamilNadu 625701, India

³Department of Biotechnology, Sethu Institute of Technology, TamilNadu 626115, India

⁴Department of International Research Center, Kalasalingam Academy of Research and Education, TamilNadu 626126, India

Received 9 July 2023; Received in revised form 24 September 2023

Accepted 9 October 2023; Available online 27 December 2023

ABSTRACT

The purpose of the current study was to determine whether the methanolic extract of *Spatoglossum asperum* has larvicidal properties against *Aedes aegypti*, *Culex quinquefasciatus*, and *Anopheles stephensi*. Various doses of methanolic seaweed extract (50, 100, 150, and 200 ppm) were applied to chosen larvae in the early 4th instar. Each test was carried out in triplicate, with a control group present simultaneously. The mortality rate was observed at regular 24 hr intervals. FTIR and GC-MS studies analyzed the methanolic extract's functional groups and phytochemical compounds. The standard plate procedure was used to test the antibacterial capacity using clinically isolated pathogens; the extract effectively killed bacteria of all types, according to findings. FTIR peaks confirmed the presence of carbonyl groups, amino acids, hydroxyl groups, and alkynes. The GC-MS study of the seaweed extract found 14 major bioactive compounds, such as citronellol and diethyl phthalate, found to have insecticidal, nematocidal, antibacterial, antioxidant, cytotoxic, antifungal, and hepatoprotective properties. DPPH radical scavenging and nitric oxide assays showed that the efficacy grew as the extract level was raised. Total phenol and flavonoid content of the methanol extract was estimated at 73.2±02 µg/mL and 89.1±0.03 µg/mL, respectively. Against 4th instar *Ae. aegypti*, *Cx. quinquefasciatus*, and *An. stephensi* larvae,

methanolic extract of *S. asperum* showed LC₅₀ values of 2.56 mg/L, 3.32 mg/L, and 2.84 mg/L, respectively. These studies suggest methanolic extracts of *S. asperum* have more potential use as a larvicidal agent against vectors of *Aedes*, *Culex*, and *Anopheles* mosquitoes.

Keywords: Antibacterial; Histopathological; Larvicidal; Methanol extract; Seaweed

1. Introduction

Mosquitoes are the most common disease-carrying insects, transmitting diseases such as malaria, chikungunya, typhoid, and dengue. The proliferation of these mosquitoes and their global distribution and the diseases they transmit significantly contributed to total mortality rates [1]. *Aedes albopictus*, *Anopheles*, *Mansonia*, and *Culex* are important disease vectors. According to the National vector control board in India, 30,627 cases of dengue fever and 15 million cases of malaria have been reported, and deaths range from 19,500 to 20,000 each year as reported by the World Health Organization [2]. Control of mosquitoes is significant in stopping the propagation of vector-borne illnesses. Although many malaria vaccines are being developed, nothing is presently accessible. Using chemical mosquito repellents causes environmental damage, and resistance can harm humans [3]. Syed et al. [4] studied the effect of kidney cell degeneration on albino mice using mosquito coils. They concluded that d-trans allethrin coming from mosquito coils is hazardous and changes creatinine and serum urea levels in 10 weeks of observation. Continuous chemical insecticides are poisonous to humans and harmful to other living organisms. Alternatives now focus on finding environmentally friendly and cost-effective mosquito repellents [5].

Humans have used plants for centuries against these deadly pathogens and other pests. Numerous studies have revealed larvicidal, pest development inhibiting, and deterrent activities of phytochemicals produced by plants [6]. The ocean contains many types of natural resources. The sea's nearly 5,000,000 lifeforms are a practically untapped source of secondary compounds [7]. As a result, experts are exploring how to kill insects

with plants and seaweed. Seaweeds are abundant in bioactive substances and compounds with therapeutic properties. They have been shown to affect insect metabolic activity through intoxication, morbidity, growth and development changes, morphological changes, oviposition, and reproduction system changes [8].

Marine algae, known as seaweeds, typically grow on the seafloor between the high and low tide marks. These algae are multicellular, polyphyletic, and oxidative. Seaweeds and other marine algae prefer rocky shores in the ocean's littoral zone for their growth [9]. More than 60 trace elements are present in marine algae at concentrations several times higher than those in higher ecosystems and these elements have shown anxiolytic and antidepressant action [10]. Elements like sodium, potassium, calcium, phosphorus, as well as vitamins are plentiful in marine algae. They contain high concentrations of the nine essential amino acids necessary for optimal wellness [11].

Spatoglossum asperum is a brown seaweed distributed all over the world. Seaweeds have been reported for their antibacterial [12-16], antioxidant [17-20], anticancer [21-25], anti-ulcer [26-30], anti-diabetic actions [31-35], and wound healing [36, 37] activities. Some seaweed-derived bioactive compounds are mosquitocidal as well [8]. *Lobophora variegata*, *Sargassum wightii*, *Ulva lactuca*, and *Caulerpa racemosa* are some species that have been reported to display anti-larvicidal activity against mosquitoes.

The brown seaweed *S. asperum* was extracted with methanol for this study to identify the extract's bioactive compounds and functional groups. By using the hole plate process, the resultant extract's antimicrobial

efficacy was examined towards microorganisms. In the larvicidal investigation, *Anopheles stephensi*, *Culex quinquefasciatus*, and *Aedes aegypti* were targeted. Changes in histology and death rates were also analyzed.




Fig. 1. Research methodology.

2. Materials and Methods

2.1 Collection and identification of seaweed

S. asperum, brown algae, was gathered from the Gulf of Mannar's Mandapam shorelines. The obtained seaweed was identified by Dr. Palanisamy, Scientist 'E' Botanical Survey of India, West Bengal. The herbarium of the collected sample is stored at Thiruvalluvar University (Fig. 1.).

2.2 Extract preparation

The gathered material was washed with sterile water and dried for 15 days in the shade. To grind the dehydrated specimens, a mechanical mixer was used, followed by a sieve. Next, 10 grams of the obtained powder were macerated with 50 mL of methanol for 48 hrs under a continuous stirrer. After that, Whatman No. 1 filter sheet was used to filter the samples. Phytochemical analysis was performed on the filtrate and its anti-bacterial and larvicidal activities were observed.

2.3 Phytochemical analysis

The goal of this common testing procedure was to detect the existence of

several botanical constituents. The methanolic extract of *S. asperum* was previously submitted for preliminary phytochemical screening [38].

2.4 FTIR analysis

The FTIR spectrum of the methanolic seaweed extract of *S. asperum* was analysed to learn more about the functional groups found in the seaweed. When undiscovered chemicals are discovered in plants, spectroscopy can be a valuable tool for elucidating their structures. The crude methanolic extract was subjected to FTIR analysis, and thus the extract was loaded in an FTIR spectroscope (FTIR-8400S, Shimadzu) with an analysis range from 500 to 4000 cm^{-1} .

2.5 GC-MS analysis

GC-MS analysis of methanolic *S. asperum* extract was performed using GCMS-QP 2010 Plus Shimadzu system. Identification and interpretation of phytochemicals on mass spectrum GC-MS were conducted using the NIST and WILEY8 Archives. The spectra of the unknown constituents were contrasted to spectra of known compounds in the NIST collection. The compounds which were spotted were given names, molecular formulas, and structures.

2.6 Total phenolic and flavonoid content

Employing standards (gallic acid and quercetin), the total phenolic content (TPC) and total flavonoid content (TFC) was established. The procedure from Singelton et al. [39], with some minor alterations, was applied to determine the TPC of various concentrations of seaweed extract (20 $\mu\text{g/mL}$ -100 $\mu\text{g/mL}$) which were made using sterile water. A mixture of 3.4 mL of 30% methanol, 0.15 mL of NaNO_2 (0.5 M), and 0.15 mL of $\text{AlCl}_3 \cdot 6\text{H}_2\text{O}$ (0.3 M) were combined with 0.3 mL of sample. After 5 minutes, 1 mL of NaOH (1 M) was added. The solution was thoroughly mixed, then absorption was read at 506 nm and compared to blank reagent. TFC was computed using a modified version of Park et

al.'s [40] formula. In a tube, 0.1 mL of test samples (20 µg/mL-100 µg/mL) were added together with 1.9 mL of distilled H₂O, 1 mL of Folin-Ciocalteu's reagent, and 1 mL of 100 g/L Na₂CO₃ to stop the process. The mixture's absorbance was gauged at 765 nm after the reaction mixture had been set aside for 2 hrs at 25°C.

2.7 Antioxidant activity

2.7.1 DPPH assay

The antioxidant capacity of extracts was estimated by UV spectrophotometer against DPPH (2,2-diphenyl-1-picrylhydrazyl) radical. For this task, the technique of Prieto, 1999 [41] was applied. In methanol, 0.1 mM solution of DPPH was added, then 1 mL of the resulting mixture was mixed with 1 mL of extract at different levels (100-1000 µg/mL). After stirring, the liquid was left alone for 30 minutes at room temperature. The absorbency was obtained at 517 nm. Ascorbic acid was used as a baseline. Higher energy is indicated by lower absorbance levels.

To determine the % of antioxidant production, the following calculation was employed.

$$\% \text{ inhibition} = \left(\frac{A_c - A_e}{A_c} \right) \times 100, \quad (2.1)$$

where A_c is the efficiency of the control and A_e is the efficiency of the tested sample.

2.7.2 Nitric oxide scavenging activity

Phosphate buffer (pH 7.4) was used to prepare 10 mM sodium nitroprusside. 1 mL of leaf extract at various concentrations was combined with 0.5 mL of sodium nitroprusside solution. The reaction mixture was incubated for 180 minutes at 25°C. Then, 0.5 mL of the incubated sample was combined with 0.5 mL of Griess reagent (1% sulphanilamide, 2% H₂PO₄, 0.1% N-1-Naphthyl Ethylenediamine Dihydrochloride). As a control, the reagent without extract was used. The absorbance was measured at 546 nm. Ascorbic acid was

used as a standard. The formula (Patel, 2010) [42] was used to figure out the % obstruction.

$$\% \text{ Scavenging activity} = \left(\frac{A_c - A_t}{A_c} \right) \times 100, \quad (2.2)$$

where A_c stands for the control's efficiency and A_t for the test's efficiency.

2.7.3 Reducing power activity

Srinivasan et al.'s [43] technique was used to find the reducing power. Several concentrations of extract (100-1000 µg/mL) were combined in 1 mL of distilled water, 2.5 mL of phosphate buffer (pH 6.6), and 2.5 mL of potassium ferricyanide (1% w/v). For 20 mins, the resulting mixture was left to stand at 50°C. After 20 mins, 2.5 mL of 20% Trichloroacetic acid was added to halt the process. For 10 mins, the mixture was mixed at 3,000 rpm. 2.5 mL of sterile water and 0.5 mL of ferric chloride (0.1%) were combined with 2.5 mL of the mixed sample, and intensity at 700 nm was obtained. Enhanced mixture absorbance suggested higher reducing power. Ascorbic acid was utilized as a reference. As a control, phosphate buffer (pH 6.6) was utilized.

2.8 Antibacterial activity of the extract

The methanolic extract of *S. asperum* was assessed for its antimicrobial efficacy against Gram-positive *Staphylococcus epidermidis*, *Bacillus subtilis*, *Micrococcus luteus*, *Streptococcus pneumoniae*, and Gram-negative *Escherichia coli*, *Klebsiella aerogenes*, *Salmonella typhimurium*, and *Vibrio parahaemolyticus*. These microbes were collected from Christian Medical College (CMC), Vellore. 20 mL of Muller Hinton Agar (MHA) (Hi-Media, Mumbai) was put into sterilized Petri dishes. Various concentrations of sample (25, 50, 75, and 100 mg/mL) were loaded in wells. Streptomycin (10 µg/disc) was used as a positive control. The loaded wells were left for 30 mins at room temperature for compound diffusion. For 24 hrs, the dishes were maintained at 37°C. The circumferences

of the zones of inhibition were assessed, and the mean values for every organism were ascertained [14]. The experiment was repeated in triplicate.

2.9 Larvicidal study

Laboratory strains of *Ae. aegypti*, *An. stephensi*, and *Cx. quinquefasciatus* were received from ICMR, VCRC, at Madurai. All procedures for larvicidal assays followed the standard WHO protocol. In a laboratory setting, the larvae were raised on plastic crates and served yeast and dog biscuits in a 1:2 ratio. Young 4th instar mosquito larvae were delivered in sets of 25 into 100 mL water-filled disposable cups. Various concentrations (50, 100, 150, and 200 ppm) of methanolic seaweed extract were introduced. Cups without any methanol extract added were used as a control. The setup was observed for 72 hrs. The mosquito death rate was recorded after 24 hrs of treatment. All experiments were done in triplicate under the precise temperature of 25°C ± 2°C and relative moisture level of 75%–85%. Dead larvae were identified by lack of movement.

2.10 Histopathological analysis

The midgut epithelial cell changes were observed after the treatment. The treatment and control of mosquito larvae of *Cx. quinquefasciatus*, *Ae. Aegypti*, and *An. Stephensi*'s 4th instars were preserved in 10% formalin. After being fixed, the samples were dehydrated and mounted in paraffin. Using a microtome, the prepared slides were sectioned. Eosin and hematoxylin were used for staining. The slides were cleaned with xylene, the pathological alterations of the sections were observed under light microscopic, and the changes were photographed. The obtained results were compared with the control group.

2.11 Statistical analysis

The information is displayed as average and standard deviation. The paired-sample t-test was implemented to establish the significance of the data. The comparisons were evaluated at various times using ANOVA with repeated measurements. We conducted our data analysis using SPSS® version 20. *P* values ≤ 0.05 were considered significant. The LC₅₀ values were measured using Probit analysis tool.


3. Results and Discussion

3.1 Phytochemical analysis

The secondary compounds found in the methanolic extract of *S. asperum* are given in Table 1. The presence of tannins, saponins, flavonoids, alkaloids, and carbohydrates in the present study of *S. asperum* is supported by several others. Andrews et al. [44] reported that the methanolic leaf extract of *S. asperum* contains tannins, phlobatannins, saponin, flavonoids, terpenoids, alkaloids, proteins, and polyphenols. *S. marginatum* has therapeutically active ingredients, as shown by the phytochemicals present in methanolic extract such as tannins, flavonoids, steroids, terpenoids, triterpenoids, and alkaloids [45]. Rajakumari and Kaleeswari [46] found substantial phytochemical substances in the methanolic extract of marine *Sargassum wightii* such as carbohydrates, proteins, phenol, saponins, glycosides, steroids, terpenoids, and alkaloids, and they concluded that it must substantiate the assertions of the health care sector and indigenous medicine that these brown seaweeds exhibit biomedically significant activities. The methanol extract of *Padina boergesenii* contains flavonoids, saponins, terpenoids, carbohydrates, phenolics, and tannins [47]. The brown seaweed *Spatoglossum squarrosum* was extracted with methanol, and phytochemical testing confirmed the presence of alkaline substances, terpenoids, polyphenolic compounds, tannins, steroids, and reducing carbohydrates [48].

Table 1. Phytochemical screening of methanolic extract of *S. asperum*.

Test	Reagents used	Result
Tannins	Ferric chloride	+
Phlobatannins	1% Aqueous HCl	-
Saponin	Distilled water	+
Flavonoids	Sodium hydroxide	+
Steroids	Phytochemical screening of methanolic extract of <i>S. asperum</i> the test steroid Salkowski test	-
Alkaloids	Dragendroff's reagent	++
Protein	Millon's reagent	-
Carbohydrate	Benedict's reagent	+
Terpenoids	Chloroform Con. Sulphuric acid	-

**Fig. 2.** FTIR investigation of methanolic extract of *S. asperum*.

3.2 FTIR analysis

FTIR outcomes revealed the occurrence of various functional groups in the methanolic extract of *S. asperum* (Fig. 2). The peak at 3524.67 cm^{-1} was assigned to OH-stretching of vibration presence of alcohols and phenols. The rise at 2885.92 cm^{-1} was given to C-H stretching of alkanes, 2976.92 cm^{-1} was given CH and CH_2 trying aliphatic group, the peak range at 1685.67 cm^{-1} -C-C indicated the presence of alkanes, the peak at 1510.16 cm^{-1} represented polyphenol skeletal structure, the peak at 1456.16 cm^{-1} indicated C-C stretching of aromatics, the peak value at 1200.61 cm^{-1} signified C-O extension of alcohols, carboxylic esters, and ethers, the range at 697 cm^{-1} assigned -C (triple bond) C-H: C-H band presence of alkynes. The methanol extract of *S. asperum* had a weak band at 3902.69 cm^{-1} , 3737.79 cm^{-1} , and 1545.81 cm^{-1} , which indicated the presence of O-H stretching of

alcohols. According to Yu et al. [8], FTIR investigations on algae and seaweed compounds indicated that the walls of algae had hazardous reaction sites of carbonyl groups, amino acids, and hydroxyl groups.

3.3 GC-MS analysis

A methanol derivative of *S. asperum*'s 14 physiologically active components have been determined by GC-MS. Retention time, molar mass, chemical formula, and biological properties of the identified compounds are given in Table 2 and Fig. 3. The most critical phytoconstituents present in the extracts were dimethyl sulfoxide (0.71), nonadecane (0.20), cyclododecane (0.21), eicosane (0.88), diethyl phthalate (4.94), pentadecanoic acid (0.59), gamolenic acid (1.67), phytol (2.44), vaccenic acid (17.03), octadecanoic acid (2.07), hexadecenoic acid (1.81), citronellal (0.65), and fucosterol (18.39). All of the bioactive

compounds were found to possess nematocidal, insecticidal, repellent, antibacterial, antifungal, cytotoxic, antioxidant properties, and properties for treating skin disorders. Citronella oil derived from citronellol is the most commonly used mosquito repellent. Diethyl phthalate and octadecanoic acid have nematocidal, insecticidal, and pesticidal activities. As reported by Manh et al. [49], the essential oil of *Cymbopogon winterianus*, also called Citronella oil, was tested against 3rd instar dengue parasite *Ae. aegypti*, and the LC₅₀

value was found to be 120.6 ppm. The toxicity effect of citronella oil was evaluated against the 3rd stage of *Ae. aegypti* by Cansian et al. [50], and the LC₅₀ value was 86.30 µg mL⁻¹; they reported that the tested samples can be used as bioinsecticides. Adsul et al. [51], derived phthalate from the extract of *Ipomoea carnea*, and the isolated compound was found to be lethal to the 4th instar stages of *Ae. aegypti* and *Cx. quinquefasciatus* with the LC₅₀ found to be 81.43 and 109.64 ppm, respectively.

Table 2. GC-MS investigation of methanolic extract of *S. asperum*.

S. No	Retention time	Compound name	Molar mass	Chemical formula	Biological properties
1	5.766	Dimethyl sulfoxide	78.13 g/mol ⁻¹	C ₂ H ₆ OS	Antibacterial, antifungal
2	11.630	Nonadecane	268.518 g/mol	C ₁₉ H ₄₀	Antimicrobial
3	12.956	Cyclododecane	168.319 g/mol	C ₁₂ H ₂₄	Antibacterial
4	14.776	Eicosane	282.5475 g/mol	C ₂₀ H ₄₂	Antifungal, antipyretic, analgesic
5	15.975	Diethyl phthalate	222.24 g/mol	C ₁₂ H ₁₄ O ₄	Antiviral, antioxidant, insecticidal, cytotoxic, antihyperglycemic, antibacterial, antifungal activity
6	17.542	Pentadecanoic acid	242.3975 g/mol	C ₁₅ H ₃₀ O ₂	Antioxidant
7	19.021	Gamolenic acid	278.43 g/mol	C ₁₈ H ₃₀ O ₂	Atherosclerosis, anti-cancer, diabetic complications, skin disorders
8	19.734	Phytol	296.53 g/mol	C ₂₀ H ₄₀ O	Antibacterial, anxiolytic, metabolism-modulating, Anticancer, radical scavenging activity, phagocytosis and cell death-inducing, antinociceptive, anti-allergic, immune-modulating, and neurotoxic effects
9	20.371	Oleic acid	282.47 g/mol	C ₁₈ H ₃₄ O ₂	Antibacterial, antiviral, antifungal, and antioxidant
10	20.371	Vaccenic acid	282.461 g/mol	C ₁₈ H ₃₄ O ₂	Antibacterial
11	20.656	Octadecanoic acid	284.48 g/mol	C ₁₈ H ₃₆ O ₂	Antioxidants, hypocholesterolemic, nematocidal, and pesticide
12	24.079	Hexadecenoic acid	254.414 g/mol	C ₁₆ H ₃₀ O ₂	Antioxidant, anti-inflammatory, hypo-cholesterolemic, and cancer prevention activities
13	25.455	Citronellol	154.25 g/mol	C ₁₀ H ₁₈ O	Insect repellent, antifungal, anti-inflammatory
14	25.455	Fucosterol	412.69 g/mol	C ₂₉ H ₄₈ O	Cytotoxic, antidiabetic, antioxidant, hepatoprotective, antihyperlipidemic, antifungal, antihistaminic, anticholinergic, antiadipogenic, anti-photodamaging, anti-osteoporotic,




Fig. 3. GC-MS investigation of a methanolic extract of *S. asperum*.

3.4 Total phenol and flavonoid content

The numerous beneficial biological impacts of flavonoids and phenolic compounds have led to substantial research on these compounds [52]. According to the quantitative examination of plant chemicals, *S. asperum* methanol extract is rich in flavonoids and phenolics. In Table 3, it is shown that the methanolic extract of *S. asperum* has an elevated phenol and flavonoid profile. Methanol's capacity to block polyphenol oxidase activity, which is responsible for the oxidation of polyphenols, makes it an ideal solvent for the extraction of polyphenolic substances [53]. The methanolic algal extract had a flavonoid content of 89.1 ± 0.03 mg QE/g and the phenol content was 73.2 ± 0.2 mg GAE/g. Tanna et al. [54], estimate the phenol and flavonoid content of the brown algae *S. asperum* and the TPC and TFC quantities were 15 ± 3 mg/mL–1, 340 ± 60 mg/mL⁻¹ they concluded that *S. asperum*, brown algae, has a higher total phenolic content and total flavonoid content than other brown marine algae. *Sargassum*, *Turbinaria* and *Padina* species were examined for their total phenolic content by Sinjel et al. [55] and the result shows that *Padina Spp* 0.135 mg GAE/g had the greatest total phenolic content, followed by *Sargassum Spp* 0.084 mg GAE/g and

Turbinaria Spp 0.004 mg GAE/g with the lowest.

Table 3. Total phenol and flavonoid content of methanolic extract of *S. asperum*.

	20	40	60	80	100
	µg/mL	µg/mL	µg/mL	µg/mL	µg/mL
Flavonoid	45.1 ± 0.03	52.1 ± 0.05	68.73 ± 0.23	77.5 ± 0.04	89.1 ± 0.03
Phenol	20.2 ± 2.9	46.0 ± 3.6	50.4 ± 0.6	66.4 ± 0.5	73.2 ± 0.2

3.5 Antioxidant activity

3.5.1 DPPH scavenging activity

DPPH assay represents one of the best ways to assess the effectiveness of antioxidants. In the current research, the *S. asperum* methanol extract was found to have potent antiradical action at therapeutic doses. Fig. 4 shows the result of methanolic extract of *S. asperum*. The antioxidant activity of *S. asperum* was lower than the standard gallic acid. A stable non-radical form of DPPH is produced when a DPPH solution is combined with a substrate that acts as a hydrogen atom donor, and the violet hue simultaneously changes to pale yellow. The methanolic extract of *S. asperum* showed the greatest efficacy of 57% at 400 µg [54]. The highest DPPH activity was $79.1 \pm 1.21\%$ in the brown seaweed *S. wightii*, which is relatively high when compared to red and green algae. Chakraborty

et al. [57] studied the DPPH activity of the brown seaweed *T. ornata* and an efficacy of 64.14% was recorded. Airanthi et al. [58] examined the DPPH activity of two brown seaweeds, *S. horneri* and *C. Hakodatensis*,

from coastal regions of Japan. The marine algae were extracted with methanol and the DPPH activity of *S. horneri* and *C. hakodatensis* were $28.50 \pm 0.75 \mu\text{g/mg}$ and $65.32 \pm 3.95 \mu\text{g/mg}$, respectively.




Fig. 4. DPPH activity of methanolic extract of *S. asperum*.

3.5.2 Nitric oxide scavenging assay

The greatest amount of suppression was seen in the methanolic extract of *S. asperum*, at 51.1 ± 0.03 . It was shown that the action was dose-dependent. Vitamin-C was used as a standard. Nitrite formation is inhibited by algae because they use oxygen for their own cellular processes instead of allowing it to interact with nitric oxide. The structure and function of numerous cellular components might be altered as a result of these chemicals. Chia YY et al. [53] evaluated the antioxidant

activity of the brown macro algae *T. ornata* and found that the methanolic extract had the most significant degree of restriction, at $54.96 \pm 0.41\%$. The brown algae *S. wightii* collected from the Mandapam coastline, was subjected to methanolic extraction and tested for its antioxidant activity. The results showed it exhibited $60.22 \pm 0.52\%$ inhibition [59]. Sodium nitroprusside (SNP) produces NO, which interacts immediately with oxygen to form nitrite. Fig. 5 shows the results of the methanol extract of *S. asperum*.




Fig.5. Nitric oxide scavenging activity of methanolic extract of *S. asperum*.

3.5.3 Reducing power activity

Reducing power is a measure of an element's ability to donate an electron, and substances with this property can serve as both

primary and secondary antioxidants by neutralizing the oxidized precursors produced during the lipid peroxidation processes [60]. Fig. 6 depicts the decreased activity of the

methanolic *S. asperum* extraction. Chandhini et al. [61] investigated the reducing power of the brown algae *Sargassum marginatum*, *Padina tetrastomatica*, and *Turbinaria conoides*. The seaweeds were extracted with methanol. With rising concentrations, the reducing power of crude methanolic extract increased. *T. conoides* and *P. tetrastomatica* have greater reducing power than the standard scavenger α -tocopherol. The methanolic extracts of *Turbinaria ornata* and *Sargassum polycystum* had high levels of reducing power.

Devi et al. [62] investigated the antioxidant abilities of green, red, and brown algae and found that among them, the brown algae *T. conoids* exhibited the highest antioxidant potential and it was lower than that of the standard. Rabecca and Doss [63] evaluated the reducing power of brown and green seaweeds and found that the methanolic extract of the brown seaweed *Lobophora variegata* displays higher reducing power compared to the green algae *Codium tomentosum*.




Fig. 6. Reducing power activity of methanolic extract of *S. asperum*.

3.6 Antibacterial activity

In this investigation, the methanolic extract of *S. asperum* showed sensible activity against both gram-negative and gram-positive bacteria. The zone of inhibition of different concentrations of extract used on the gram-negative and gram-positive bacteria is given in Tables 4-5 below. In gram-negative microbes, the solvent has maximum activity against nosocomial and pathogenic *K. aerogenes* 16 ± 1.7 mm and gram-positive pneumococcal pathogen *S. pneumoniae* 18 ± 1.0 mm. Hasan et al. [64] measured the antibacterial activity of the brown seaweeds *Turbinaria conoides*, *Padina gymnospora*, and *Sargassum tenerimum*. They found that the methanol extract of all the selected seaweed had moderate activity against *E. coli*, *S. epidermidis*, *V. Parahemolyticus*, and *B. Subtilis*. Further, they concluded that the strongest antibacterial activity was displayed by the methanol extract among seven solvents used in their study. Taskin et al. [65] investigated the antimicrobial properties of

Cystoseira barbata, against gram-negative *E. coli*, *E. aerogenes* and gram-positive *Micrococcus luteus* and found it exhibited the highest activity against the tested pathogens. The crude extract of *Gracilaria edulis* inhibits the progress of *E. aerogenes* and *E. coli* [66]. The methanolic extract of *Padina australis* shows maximum inhibition towards gram-positive *B. subtilis*, *S. aureus* and less activity towards gram-negative *E. coli*. According to the current study's outcomes, gram-positive bacteria responded to the extracts more readily than gram-negative bacteria did. A possible explanation is a hydrophobic lipopolysaccharide found in the outer membrane of gram-negative bacteria, which defends against many chemicals by possibly preventing the active compounds from entering the cell [67]. *Sargassum latifolium* and *Sargassum platicarpum* were extracted with methanol and were shown to have good action towards *S. xylosus*, *E. Coli*, and *Salmonella Spp* [68]. Pandithurai et al. [69] reported that the methanolic extract of *S. asperum* has moderate

action against *S. aureus* and *K. pneumoniae*. Two seaweed species, *Hypnea musciformis* and *Enteromorpha intestinalis* were tested against human pathogens *Klebsiella Spp* and *S. aureus* and compared with gentamycin. Methanol has consistently been the greatest active solvent for retrieving antimicrobial properties from the chosen seaweeds. *Sargassum vulgare* and *Sargassum fusiforme* were tested against multidrug-resistant clinically isolated gram-negative and gram-positive bacteria. The extract showed

moderate activity against *S. aureus* and *K. pneumoniae*. Morphological changes in bacterial strains have been observed through transmission electron microscopy, indicating diminished cell size and cell wall separation [70]. The methanol-based extracts of *Sargassum wightii* and *Stechospermum marginatum* were efficient against *E. coli*, *E. faecalis*, *S. epidermidis*, *M. luteus*, *P. aeruginosa*, *S. aureus*, and the human fungus *Candida albicans*, as determined by the well plate technique [71].

Table 4. Antimicrobial action of methanolic crude extract against gram-negative bacteria.

Pathogens	Antibiotic	Extract without methanol	25 (mg/mL)	50 (mg/mL)	75 (mg/mL)	100 (mg/mL)
<i>Escherichia coli</i>	20±1.5	17±1.2	15±1.5	16±1.1	18±1.8	19±1.1
<i>Klebsiella aerogenes</i>	22±1.5	15±1.8	16±1.7	16±1.8	17±1.5	19±2.0
<i>Salmonella typhimurium</i>	20±1.5	16±0.9	14±1.5	15±2.0	18±2.5	18±1.5
<i>Vibrio parahaemolyticus</i>	17±1.5	12±1.5	13±1.1	14±0.5	15±0.5	16±0.5

Table 5. Antimicrobial activity of methanolic crude extract against gram-positive bacteria.

Pathogens	Antibiotic	Extract without methanol	25 (mg/mL)	50 (mg/mL)	75 (mg/mL)	100 (mg/mL)
<i>Staphylococcus epidermidis</i>	21±2.0	11±1.8	14±1.5	14±1.0	15±2.5	15±2
<i>Bacillus subtilis</i>	19±1.5	14±1.7	12±0.5	13±0.5	15±1.0	16±1.0
<i>Micrococcus luteus</i>	22±0.2	17±2.8	16±1.5	18±1.5	19±2.0	20±3.1
<i>Streptococcus pneumoniae</i>	21±2.0	16±1.9	18±1.0	18±2.0	19±2.0	21±3.3

Results are represented as average± SD, with results representing the mean of trials carried out in triplicate. The zone of inhibition was measured in mm.

3.7 Larvicidal activity

Tables 6-7 display the insecticidal efficacy of *S. asperum* methanolic extract against *Aedes aegypti*, *Culex quinquefasciatus*, and *Anopheles stephensi*. Of the three species of mosquitoes studied, only *Ae. aegypti* was significantly affected by the extract. The mortality rate was *Ae. aegypti* (2.56 mg/L), *An. stephensi* (2.84 mg/L), and *Cx. quinquefasciatus* (3.32 mg/L), respectively at the concentration of 200 ppm after 72 hrs of action. The larvicidal activity of *Sargassum microcystum* against 4th instar larvae of *Ae. aegypti*, *C. quinquefasciatus*, and *An. Stephensi* was studied. It was concluded that the activity may be due to the occurrence of saponins and triterpenoids in the product [6]. The methanolic extract of *S. asperum*, *S. marginatum*, and *S. wightii* were tested against the 2nd and 3rd instars of *Cx. quinquefasciatus* and *Ae. aegypti* and after 24 hrs, 100% mortality was recorded. The activity was

higher in 2nd instar specimens compared with 3rd instar specimens [72]. Hira et al. [73] studied the larvicidal potential of brown seaweeds *S. ilicifolium*, *S. binderi*, *S. lanceolatum*, and *S. variabile* against 4th instar *Ae. aegypti*. Among them, *S. binderi* showed a 50% mortality at low concentration. The methanol extract of *Sargassum binderi* caused larvicidal and histological changes in *Aedes Aegypti* with an LC₅₀ value of 192.43 g/mL [8]. Extracts from *S. wightii*, *S. ilicifolium*, and *G. acerosa* have been studied for their larvicidal efficacy towards 4th instar *Ae. aegypti*, *Cx. quinquefasciatus*, and *An. stephensi*. They explained the way seaweed extracts were tested on *Daphnia* and how they might have possessed crucial bioactive chemicals that led to the high larval mortality. The outcome shows that it is non-toxic to non-target organisms [74]. *Turbinaria conoids* extract was tested against *Cx. quinquefasciatus* and 80% mortality was observed after 24 hrs of

incubation [75]. The methanol extract of *G. corticate* was tested against the malarial vector

An. stephensi and the LC₅₀ value was 498.91 mg/L [6].

Table 6. Mortality rate of selected mosquito species after being exposed to various concentrations of *S. asperum* methanolic extract for 72 hours.

Conc. of <i>S. asperum</i> methanol extract	No. of larvae per application (N)	Number of larval mortalities		
		<i>Ae. aegypti</i>	<i>Cx. quinquefasciatus</i>	<i>An. stephensi</i>
Control	25	0	0	0
50 ppm	25	4	2	5
100 ppm	25	8	7	6
150 ppm	25	14	8	8
200 ppm	25	19	14	13

Table 7. Larvicidal effect of methanolic extract of *S. asperum* against 4th instar larva of *An. stephensi*, *Cx. quinquefasciatus*, *Ae. aegypti*.

Name of the species	LC ₅₀ (mg/L)	Regression equation	R ²
<i>Aedes aegypti</i>	2.56	$Y=3.5x-4.7$	0.9511
<i>Culex quinquefasciatus</i>	3.32	$Y=3.4x-4$	0.957
<i>Anopheles stephensi</i>	2.854	$Y=2.9x-2.3$	0.9428

3.8 Histopathological analysis

The midgut of *An. stephensi*, *Ae. Aegypti*, and *Cx. quinquefasciatus* larvae was observed after being exposed to methanolic *S. asperum* extract at various concentrations. Histopathological analysis indicated that the damage was in the midgut, particularly in the corneal epithelium, microvilli, and basal membrane. The histological variations in both untreated and treated larval mosquitoes are shown in Figs. 7-9. Conferring with Rohmah et al. [76], the midgut of larvae was damaged primarily due to several roles occurring in this

location, such as the digestive process, nutrition captivation, signal transduction, and osmoregulation. This was triggered by the extract's deadly effect on the columnar epithelium layering the midgut. This disrupted enzyme excretion and the digestive process, leaving larvae with an energy deficit that could lead to death [77]. At 200 ppm, the extract causes severe damage to the peritrophic membrane and epithelial cells. The changes were due to the presence of larvicidal compounds such as citronella and dimethyl phthalate in the extract. According to a report by Yu et al. [8], the occurrence of vesicles of differing shapes, damage to microvilli, and inflammation of cells were observed in the anterior midgut epithelial cells and the posterior intestinal lumen epithelial of control larvae. On the other hand, the epithelium was lined by huge, irregular cells with massive globular bases were observed *Ae. aegypti* larvae treated with *Sargassum binderi* extract.




Fig. 7. Histopathological changes in *An. stephensi* treated with methanolic extract of *S. asperum* [A] Control larvae, [B] Treated larvae, EP (Epithelial cells), BM (Basal membrane), PM (Peritrophic membrane), LC (cell lysis).




Fig. 8. Histopathological changes in *Ae. aegypti* treated with methanolic extract of *S. asperum* [A] Control larvae, [B] Treated larvae, EP (Epithelial cells), BM (Basal membrane), PM (Peritrophic membrane), LC (cell lysis).




Fig. 9. Histopathological changes in *Cx. quinquefasciatus* treated with methanolic extract of *S. asperum* [A] Control larvae, [B] Treated larvae, EP (Epithelial cells), BM (Basal membrane), PM (Peritrophic membrane), DC (Destroyed cell), BE (Elongated Brush border).

4. Conclusion

The efficacy of bioactive substances against various insect pests, such as mosquitoes, has been the subject of numerous studies in recent decades. This investigation concluded that methanolic extract of *S. asperum* induced noticeable larval death and showed moderate activity against all the tested organisms. The extract also has good activity against all the verified clinical pathogens. The GC-MS analysis identified 14 biologically active compounds. The methanolic extract of *S. asperum* contains a high quantity of phenol and flavonoids. The DPPH, reducing power, and nitric oxide scavenging assay results from this study are in agreement with the extracts' antioxidant activity shown in other studies. The plant's larvicidal, antibacterial, and

antioxidant activities are generally supported by the research. At 200 ppm, methanolic extract of *S. asperum* has high activity against all the tested larvae. After 72 hrs exposure, the highest activity was recorded against *Ae. aegypti* and the LC_{50} rate was 2.56 mg/L. Histopathological damage occurred in the treated groups' peritrophic membrane, cell wall, and epithelial cells compared to control larvae. In accordance with the study's findings, this seaweed could be useful in the development of innovative mosquitocidal formulations in the future and may have beneficial medical applications. Further research is necessary to validate the larvicidal activity, antibacterial, and antioxidant effects, as well as to identify and isolate the active chemicals, and clarify their pharmacological characteristics. This will

include performing in vitro and in vivo studies to understand their antibacterial and antioxidant effects, and to find and isolate the active chemicals, and clarify their pharmacological characteristics. As a result, additional compound fractionation and purification will clarify which compounds are responsible for these effects.

Acknowledgements

The authors of this article would like to express their gratitude to Thiruvalluvar University, Serkkadu, Vellore-632 115. and also thankful to ICMR, Vector Control Research Centre, Madurai to provide the mosquito eggs.

References

- [1] Kaura T, Walter NS, Kaur U, Sehgal R. Different Strategies for Mosquito Control: Challenges and Alternatives. In Mosquito Research-Recent Advances in Pathogen Interactions, Immunity, and Vector Control Strategies. IntechOpen. 2022 Apr 17.
- [2] Malaria-World Health Organization (WHO). Available from: <https://www.who.int>.
- [3] Roy DN, Goswami R, Pal A. The insect repellents: A silent environmental chemical toxicant to the health. Environmental toxicology and pharmacology, 2017;50:91-102.
- [4] Syed M, Tousia S, Balowria KS, Jabeen N, Faruqi NA, Beg A. Haematological and Histopathological Effects of Mosquito Coil Smoke on Kidney-An Experimental study on Albino Rat. Acedamia Anatomica International, 2019;5(1):49-53.
- [5] Maia MF, Moore SJ. Plant-based insect repellents: a review of their efficacy, development and testing. Malaria journal, 2011;10(1):1-5.
- [6] Ali MY, Ravikumar S, Beula JM. Mosquito larvicidal activity of seaweeds extracts against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*. Asian Pacific Journal of Tropical Disease, 2013;3(3):196-201.
- [7] Balaraman P, Balasubramanian B, Liu WC, Kaliannan D, Durai M, Kamyab H, Alwetaishi M, Maluventhen V, Ashokkumar V, Chelliapan S, Maruthupandian A. Sargassum myriocystum-mediated TiO₂-nanoparticles and their antimicrobial, larvicidal activities and enhanced photocatalytic degradation of various dyes. Environmental research, 2022;204:112278.
- [8] Yu KX, Wong CL, Ahmad R, Jantan I. Larvicidal activity, inhibition effect on development, histopathological alteration and morphological aberration induced by seaweed extracts in *Aedes aegypti* (Diptera: Culicidae). Asian Pacific journal of tropical medicine, 2015;8(12):1006-12.
- [9] Doss A, Rukshana MS. Distribution pattern of marine seaweeds in the Manapad coastal region. Saudi J. Pathol. Microbiol, 2016;1(1):10-3.
- [10] Adeola AO, Ore OT, Fapohunda O, Adewole AH, Akerele DD, Akingboye AS, Oloye FF. Psychotropic drugs of emerging concerns in aquatic systems: ecotoxicology and remediation approaches. Chemistry Africa, 2022;5(3):481-508.
- [11] Godswill AG, Somtochukwu IV, Ikechukwu AO, Kate EC. Health benefits of micronutrients (vitamins and minerals) and their associated deficiency diseases: A systematic review. International Journal of Food Sciences, 2020;3(1):1-32.
- [12] Martín-Martín RP, Carcedo-Forés M, Camacho-Bolós P, García-Aljaro C, Angulo-Preckler C, Avila C, Lluch JR, Garreta AG. Experimental evidence of antimicrobial activity in Antarctic seaweeds: ecological role and antibiotic potential. Polar Biology, 2022;45(5):923-36.

- [13] Mofeed J, Deyab M, Mohamed A, Moustafa M, Negm S, El-Bilawy E. Antimicrobial activities of three seaweeds extract against some human viral and bacterial pathogens. *Biocell*, 2022;46(1):247.
- [14] Alghazeer R, Whida F, Abduelrhman E, Gammoudi F, Azwai S. Screening of antibacterial activity in marine green, red and brown macroalgae from the western coast of Libya. *Natural sciences*, 2013;5(1):26511.
- [15] Charway GN, Yenumula P, Kim YM. Marine algae and their potential application as antimicrobial agents. *Journal of Food Hygiene and Safety*, 2018;33(3):151-6.
- [16] Pérez MJ, Falqué E, Domínguez H. Antimicrobial action of compounds from marine seaweed. *Marine drugs*, 2016;14(3):52.
- [17] Rajeshkumar S, Parameswari RP, Jayapriya J, Tharani M, Ali H, Aljarba NH, Alkahtani S, Alarifi S. Apoptotic and antioxidant activity of gold nanoparticles synthesized using marine brown seaweed: An in vitro study. *BioMed Research International*, 2022.
- [18] Michalak I, Tiwari R, Dhawan M, Alagawany M, Farag MR, Sharun K, Emran TB, Dhama K. Antioxidant effects of seaweeds and their active compounds on animal health and production—a review. *Veterinary Quarterly*, 2022;42(1):48-67.
- [19] Chew YL, Lim YY, Omar M, Khoo KS. Antioxidant activity of three edible seaweeds from two areas in South East Asia. *LWT-Food Science and Technology*, 2008;41(6):1067-72.
- [20] Pinteus S, Silva J, Alves C, Horta A, Fino N, Rodrigues AI, Mendes S, Pedrosa R. Cytoprotective effect of seaweeds with high antioxidant activity from the Peniche coast (Portugal). *Food chemistry*, 2017;218:591-9.
- [21] Figueroa FA, Abdala-Díaz RT, Pérez C, Casas-Arrojo V, Nesic A, Tapia C, Durán C, Valdes O, Parra C, Bravo-Arrepol G, Soto L. Sulfated polysaccharide extracted from the green algae *Codium bernabei*: Physicochemical characterization and antioxidant, anticoagulant and antitumor activity. *Marine Drugs*, 2022;20(7):458.
- [22] Viswanathan S, Palaniyandi T, Kannaki P, Shanmugam R, Baskar G, Rahaman AM, Paul LT, Rajendran BK, Sivaji A. Biogenic synthesis of gold nanoparticles using red seaweed *Champia parvula* and its anti-oxidant and anticarcinogenic activity on lung cancer. *Particulate Science and Technology*, 2023;41(2):241-9.
- [23] Gutiérrez-Rodríguez AG, Juárez-Portilla C, Olivares-Bañuelos T, Zepeda RC. Anticancer activity of seaweeds. *Drug Discovery Today*, 2018;23(2):434-47.
- [24] Rocha DH, Seca AM, Pinto DC. Seaweed secondary metabolites in vitro and in vivo anticancer activity. *Marine drugs*, 2018;16(11):410.
- [25] Karkhane Yousefi M, Seyed Hashtroudi M, Mashinchian Moradi A, Ghasempour AR. In vitro investigating of anticancer activity of focuxanthin from marine brown seaweed species. *Global Journal of Environmental Science and Management*, 2018;4(1):81-90.
- [26] El-Bilawy EH, Al-Mansori AN, Soliman SA, Alotibi FO, Al-Askar AA, Arishi AA, Sabry AE, Elsharkawy MM, Heflish AA, Behiry SI, Abdelkhalek A. Antifungal, Antiviral, and HPLC Analysis of Phenolic and Flavonoid Compounds of *Amphiroa anceps* Extract. *Sustainability*, 2022;14(19):12253.
- [27] Remya RR, Julius A, Ramadoss R, Parthiban S, Bharath N, Pavana B, Samrot AV, Kanwal S, Vinayagam M, Gameda FW. Pharmacological Activities of Natural Products from Marine Seaweed

- Turbinaria ornata: A Review. Journal of Nanomaterials, 2022 Sep 8.
- [28] Senthil KA, Murugan A. Antiulcer, wound healing and hepatoprotective activities of the seaweeds Gracilaria crassa, Turbinaria ornata and Laurencia papillosa from the southeast coast of India. Brazilian Journal of Pharmaceutical Sciences, 2013; 49:669-78.
- [29] Liu L, Heinrich M, Myers S, Dworjanyn SA. Towards a better understanding of medicinal uses of the brown seaweed Sargassum in Traditional Chinese Medicine: A phytochemical and pharmacological review. Journal of ethnopharmacology, 2012;142(3):591-619.
- [30] Devi JA, Balan GS, Periyannayagam K. Pharmacognostical study and phytochemical evaluation of brown seaweed Sargassum wightii. Journal of Coastal Life Medicine, 2013;1(3):199-204.
- [31] Siddiqui NZ, Rehman AU, Yousuf W, Khan AI, Farooqui NA, Zang S, Xin Y, Wang L. Effect of crude polysaccharide from seaweed, Dictyopteris divaricata (CDDP) on gut microbiota restoration and anti-diabetic activity in streptozotocin (STZ)-induced T1DM mice. Gut Pathogens, 2022;14(1):39.
- [32] Nadeeshani H, Hassouna A, Lu J. Proteins extracted from seaweed Undaria pinnatifida and their potential uses as foods and nutraceuticals. Critical Reviews in Food Science and Nutrition, 2022;62(22):6187-203.
- [33] Senthil KA, Murugan A. Antiulcer, wound healing and hepatoprotective activities of the seaweeds Gracilaria crassa, Turbinaria ornata and Laurencia papillosa from the southeast coast of India. Brazilian Journal of Pharmaceutical Sciences, 2013;49:669-78.
- [34] Pati MP, Sharma SD, Nayak LA, Panda CR. Uses of seaweed and its application to human welfare: A review. Int. J. Pharm. Pharm. Sci, 2016; 8:12-20.
- [35] Bhatia S, Sharma K, Sharma A, Nagpal K, Bera T. Anti-inflammatory, Analgesic and Antiulcer properties of Porphyra vietnamensis. Avicenna journal of phytomedicine, 2015;5(1):69.
- [36] Shen S, Chen X, Shen Z, Chen H. Marine polysaccharides for wound dressings application: An overview. Pharmaceutics, 2021;13(10):1666.
- [37] Mateos R, Pérez-Correa JR, Domínguez H. Bioactive properties of marine phenolics. Marine drugs, 2020;18(10):501.
- [38] Harborne AJ. Phytochemical methods a guide to modern techniques of plant analysis. springer science & business media; 1998 Apr 30.
- [39] Singleton VL, Orthofer R, Lamuela-Raventós RM. [14] Analysis of total phenols and other oxidation substrates and antioxidants by means of folin-ciocalteu reagent. InMethods in enzymology, Academic press, 1999;299 152-78.
- [40] Park YS, Jung ST, Kang SG, Heo BG, Arancibia-Avila P, Toledo F, Drzewiecki J, Namiesnik J, Gorinstein S. Antioxidants and proteins in ethylene-treated kiwifruits. Food Chemistry, 2008;107(2):640-8.
- [41] Prieto P, Pineda M, Aguilar M. Spectrophotometric quantitation of antioxidant capacity through the formation of a phosphomolybdenum complex: specific application to the determination of vitamin E. Analytical biochemistry, 1999;269(2):337-41.
- [42] Patel A, Patel A, Patel NM. Determination of polyphenols and free radical scavenging activity of Tephrosia purpurea linn leaves (Leguminosae). Pharmacognosy Research, 2010;2(3):152-8.
- [43] Srinivasan R, Ravali B, Suvarchala P, Honey A, Tejaswini A, Neeraja P. Leucas

- aspera-medicinal plant: a review. International Journal of Pharma and Bio Sciences, 2011;2(1):153-9.
- [44] Andrews S, Azhagu Madhavan S, Ganesan S, Arjun P, Jeyaprakash R, Baskara Sanjeevi S, Ramasamy M. Different bioactive constituents and biochemical composition of brown seaweed *Spatoglossum marginatum*. Waffnen-Und Kostumkunde Journal. 2020.
- [45] Madhavan SA. Different bioactive constituents and biochemical composition of brown seaweed *spatoglossum marginatum*, 2021;2020;7(4):349-56.
- [46] Rajakumari M, Kaleeswari B. Phytochemical analysis of marine brown algae (*Sargassum wightii*) from gulf of mannar, mandabam, India. IMIRJ, III, 2019;(2):2456-4613.
- [47] Ragunath C, Kumar YAS, Kanivalan I, Radhakrishnan S. Phytochemical screening and gc-ms analysis of bioactive constituents in the methanolic extract of *Caulerpa racemosa* (Forssk.) J. Agardh and *Padina boergesenii* Allender & kraft. *Current Applied Science and Technology*, 2020, pp. 380-93.
- [48] Sekar A, Sudha TPS, Sivalingam AM, Sivamani G, Pandian A. Effective bioactive compounds and fluorescence analysis of brown seaweed *Spatoglossum asperum* J. Agardh. Upjoz. 2022;43(24):493-9. Available From: <https://www.mbmph.com/index.php/upjoz/article/view/3351>.
- [49] Manh HD, Hue DT, Hieu NT, Tuyen DT, Tuyet OT. The Mosquito larvicidal activity of essential oils from *Cymbopogon* and *Eucalyptus* Species in Vietnam. *Insects*, 2020;11(2):128.
- [50] Cansian RL, Staudt A, Bernardi JL, Puton BM, Oliveira D, de Oliveira JV, Gomes AC, Leal IC, Simas NK, Zeni J, Jungues A. Toxicity and larvicidal activity on *Aedes aegypti* of citronella essential oil submitted to enzymatic esterification. *Brazilian Journal of Biology*, 2021, pp. 83.
- [51] Adsul VB, Khatiwora E, Torane RC, Deshpande NR. Isolation and characterization of dibutyl phthalate from leaves of *Ipomoea carnea*. *Chemistry of Natural Compounds*, 2012;48:7123.
- [52] Generalić Mekinić I, Skroza D, Šimat V, Hamed I, Čagalj M, Popović Perković Z. Phenolic content of brown algae (*Pheophyceae*) species: Extraction, identification, and quantification. *Biomolecules*, 2019;9(6):244.
- [53] Chia YY, Kanthimathi MS, Khoo KS, Rajarajeswaran J, Cheng HM, Yap WS. Antioxidant and cytotoxic activities of three species of tropical seaweeds. *BMC Complementary and Alternative Medicine*, 2015;15(1):1-4.
- [54] Tanna B, Choudhary B, Mishra A, Chauhan O.P, Patel MK, Shokralla S, El-Abedin TKZ, Elansary HO, Mahmoud EA. Antioxidant, scavenging, reducing, and anti-proliferative activities of selected tropical brown seaweeds confirm the nutraceutical potential of *Spatoglossum asperum*. *Foods*, 2021;10(10):2482.
- [55] Sinjal CA, Rompas RM, Sumilat DA, Suryanto E. Antioxidant and photoprotective activity of brown seaweed from North Sulawesi Coast. *International Journal of ChemTech Research*, 2018; 11:121-33.
- [56] Siddhartha AG, Gupta VK. Green Synthesis Of MgO Nanoparticles Prepared By *Ficus religiosa* And Monitoring of Their Antimicrobial Activity Against *Pseudomonas aeruginosa*. *Solid State Technology*, 2020;63(6):3259-66.
- [57] Chakraborty K, Praveen NK, Vijayan KK, Rao GS. Evaluation of phenolic contents and antioxidant activities of brown seaweeds belonging to *Turbinaria* spp.(*Phaeophyta*, *Sargassaceae*) collected from Gulf of Mannar. *Asian Pacific*

- journal of tropical biomedicine, 2013;3(1):8-16.
- [58] Airanthi MW, Hosokawa M, Miyashita K. Comparative antioxidant activity of edible Japanese brown seaweeds. Journal of Food Science, 2011;76(1):C104-11.
- [59] Indu H, Seenivasan R. In vitro antioxidant activity of selected seaweeds from southeast coast of India. International Journal of Pharmacy and Pharmaceutical Sciences, 2013;5(2):474-84.
- [60] Kurutas EB. The importance of antioxidants which play the role in cellular response against oxidative/nitrosative stress: current state. Nutrition journal, 2015;15(1):1-22.
- [61] Chandini SK, Ganesan P, Bhaskar N. In vitro antioxidant activities of three selected brown seaweeds of India. Food chemistry, 2008;107(2):707-13.
- [62] Devi GK, Manivannan K, Thirumaran G, Rajathi FA, Anantharaman P. In vitro antioxidant activities of selected seaweeds from Southeast coast of India. Asian Pacific journal of tropical medicine, 2011;4(3):205-11.
- [63] Rebacca R, Doss A, A comparative study on the antioxidant activity of extracts of *Lobophora variegata* Less and *Codium tomentosum* STACKHOUSE. IJBPAS, 2022;11(5):2382-96.
- [64] Hasan M, Imran MA, Bhuiyan FR, Ahmed SR, Shanzana P, Moli MA, Foysal SH, Dabi SB. Phytochemical constituency profiling and antimicrobial activity screening of seaweeds extracts collected from the Bay of Bengal seacoasts. bioRxiv. 2019:680348.
- [65] Taskin E, Ozturk M, Kurt O. Antibacterial activities of some marine algae from the Aegean Sea (Turkey). African journal of Biotechnology, 2007;6(24):2746.
- [66] Mahyoub JA. Mosquito Larvicidal activity of seaweed extracts against *Anopheles d'thali* with reference to its side effects on aquatic nontraget organisms. International Journal of Mosquito Research, 2018;5(6):34-8.
- [67] Klimjit A, Praiboon J, Tiengrim S, Chirapart A, Thamlikitkul V. Phytochemical composition and antibacterial activity of brown seaweed, *Padina australis* against human pathogenic bacteria. Journal of Fisheries and Environment, 2021;45(1):8-22.
- [68] Moubayed NM, Al Hourri HJ, Al Khulaifi MM, Al Farraj DA. Antimicrobial, antioxidant properties and chemical composition of seaweeds collected from Saudi Arabia (Red Sea and Arabian Gulf). Saudi journal of biological sciences, 2017;24(1):162-9.
- [69] Pandithurai M, Murugesan S, Sivamurugan V. Antibacterial activity of various solvent extracts of marine brown alga *Spatoglossum asperum*. International Journal of Pharmacological Research, 2015;5(6):133-8.
- [70] El Shafay SM, Ali SS, El-Sheekh MM. Antimicrobial activity of some seaweeds species from Red sea, against multidrug resistant bacteria. The Egyptian Journal of Aquatic Research, 2016;42(1):65-74.
- [71] Shanmughapriya S, Manilal A, Sujith S, Selvin J, Kiran GS, Natarajaseenivasan K. Antimicrobial activity of seaweeds extracts against multiresistant pathogens. Annals of Microbiology, 2008; 58:535-41.
- [72] Kumar RS, Sujith S. In vitro mosquito larvicidal activity of marine algae against the human vectors, *Culex quinquefasciatus* (Say) and *Aedes aegypti* (Linnaeus) (Diptera: Culicidae). International Journal of Zoological Research, 2011;7(3):272-8.
- [73] Hira K, Tariq RM, Sultana V, Ara J, Ehteshamul-Haque S. Effect of seaweeds occurring at Karachi coast on mosquito larvae and liver function in rats. Pak. J. Pharm. Sci, 2017;30(2):387-91.

- [74] Perumal P, Dhanasundaram S, Aravinth A, Amutha V, Santhanam P. Larvicidal property of the extracts of the seaweeds; *Sargassum wightii*, *S. ilicifolium* and *Gelidiella acerosa* against *Anopheles stephensi*, *Aedes aegypti* and *Culex quinquefasciatus*. *Biocatalysis and Agricultural Biotechnology*, 2022;43:102436.
- [75] Kumari K, Suresh A, Gallyot EJ, Gurunadan V, Velmurugan N. Mosquito larvicidal and bactericidal action of crude extract of marine brown alga *Turbinaria conoides*. *Asia-Pacific J Sci Technol*, 2020;25:1-9.
- [76] Rohmah EA, Subekti S, Rudyanto M. Larvicidal Activity and Histopathological Effect of *Averrhoa bilimbi* Fruit Extract on *Aedes aegypti* from Surabaya, Indonesia. *Journal of Parasitology Research*. 2020;2020.
- [77] Pavananundt P, Jiraungkoorskul K, Kosai P, Jiraungkoorskul W. Larvicidal properties of *Cassia siamea* leaf against *Aedes aegypti* larvae. *International Journal of Modern Agriculture*, 2013;2(1):1-8.