

Species Composition and Spatial Distribution of Dominant Trees in the Forest Ecotone of a Mountain Ecosystem, Northern Thailand

Dokrak Marod^{1,2}, Sutteera Hermhuk^{1,3*}, Sarawood Sungkaew^{1,2},
Sathid Thinkampheang⁴, Torlarp Kamy⁵ and Wimonmart Nuipakdee⁶

¹Department of Forest Biology, Faculty of Forestry, Kasetsart University, Bangkok 10900, Thailand

²Center for Advanced Studies in Tropical Natural Resources, NRU-KU, Kasetsart University, Bangkok 10900, Thailand

³Faculty of Agricultural Production, Maejo University, Chiang Mai 50290, Thailand

⁴Cooperation Centre of Thai Forest Ecological Research Network, Kasetsart University, Bangkok 10900, Thailand

⁵Department of Agroforestry, Maejo University, Phrae Campus, Phrae 54140, Thailand

⁶Man and Biosphere Programme, Mae Sa-Kog Ma Biosphere Reserve Office, Department of National Parks, Wildlife and Plant Conservation, Bangkok 10900, Thailand

ARTICLE INFO

Received: 9 Oct 2018
Received in revised:
11 Jan 2019
Accepted: 29 Jan 2019
Published online:
19 Mar 2019
DOI: 10.32526/ennrj.17.3.2019.21

Keywords:

Doi Suthep-Pui National Park/
Ecological niche/ Forest
ecotone/ Long-term ecological
research /Permanent plot

* Corresponding author:

E-mail: h.sutteera@gmail.com

ABSTRACT

Plants' ecological niches are important to study, particularly for applying the knowledge to restoration programs. This study clarified the relationships of tree species composition and spatial distribution to environmental factors in a mountain ecosystem. A 3 ha permanent plot was established across the forest ecotone at Doi Suthep-Pui National Park, Chiang Mai Province, Thailand. The spatial distributions of the 20 dominant tree species were analyzed using a generalized linear model (GLM) combined with geographic information system (GIS) techniques. High species diversity was found, including 165 species representing 118 genera and 59 families. Elevation strongly influenced tree distribution, which could be divided into three groups: low-altitude species in deciduous dipterocarp forest (DDF), high-altitude species in lower montane forest (LMF) and coexisting DDF and LMF species. The GLM analysis revealed that soil texture, which ranged from sandy to clayey, influenced tree distribution. The results suggest that restoration programs should select suitable species based on their niches.

1. INTRODUCTION

Environmental factors control plant regeneration processes, including plant establishment, reproduction and distribution (Guisan and Zimmermann, 2000; Turner, 2001). Many researchers have defined two types of ecological gradient affecting plant distributions. The first includes direct factors that govern physiological processes (nutrients and climatic factors) and the second comprises indirect factors (such as topographic factors, habitat type and geology) that have no direct physiological relevance to species performance or survival strategies (Guisan et al., 1999; Austin, 2002; Bacaro et al., 2008). Indirect factors can be measured in the field and are often used because they are correlated with the observed tree spatial distributions.

Studies based on long-term ecological research have reported that large permanent plots can be used to predict plant distribution (Bunyavejchewin et al.,

2003; Valencia et al., 2004). Such plots not only provide information about the forest dynamics but can also be used to study species distribution (Dallmeier, 1992; Condit et al., 1999). Statistical approaches such as canonical correspondence analysis or logistic regression are used to analyze species distribution as a function of environmental factors (Guisan and Zimmermann, 2000; Lan et al., 2011; Asanok et al., 2017). For instance, the dominant species is determined based on the total number of trees (Li et al., 2009; Lan et al., 2011; Asanok et al., 2017; Walthert and Meier, 2017) and quantitative values such as the importance value index (Campos et al., 2000) and total basal area (Homeier et al., 2010; Höfle et al., 2014) are used as dependent variables. The results reflect the appearance of a niche or the suitability of a species in a particular area (Lan et al., 2011). A tree distribution map often needs to be coupled with geographic information systems (GIS), which have become popular in recent forest ecology

studies (Caillaud et al., 2010; Walthert and Meier, 2017), particularly in tropical forests, where high plant species diversity is associated with environmental conditions (Condit et al., 1999; Harms et al., 2001; Lan et al., 2011). Although studies have covered almost all forest types, few have documented the forest ecotone (FE) between lowland and upland forest along altitudinal gradients.

The mountain ecosystem in the upper reaches of the Ping River at Doi Suthep-Pui, Chiang Mai Province, Northern Thailand, is an important area for plant conservation (Fukushima et al., 2008). The environmental factors responsible for plant distribution can vary in the FE located between lowland and upland forest (Tang and Ohsawa, 1997), especially the topographic and climatic factors (Martin et al., 2007). In addition, soil properties in this area also vary due to the soil parent material, which is related to topographical and geological gradients (Rhodes et al., 2005; Weill and Munkholm, 2015) or vegetation cover (Ueda et al., 2017). Deciduous and evergreen species coexist in the FE located at altitudes between deciduous dipterocarp forest (DDF) and lower montane forest (LMF) (Marod et al., 2015). The plants in this FE are very sensitive to environmental changes (Odum, 1959; Frelich, 2002; Marchand and Houle, 2006), especially increases in temperature and drought (Ameztegui and Coll, 2011). Marod et al. (2015) reported that deciduous species, which mostly occupy the DDF and are highly adapted to drought, can occur at higher altitudes. By contrast, evergreen species are susceptible to environmental changes. Therefore, rapid environmental changes may influence their regeneration and distribution (Ruiz et al., 2008). However, little long-term ecological research has examined the adaptation and migration of tree species in a LMF in relation to changes in environmental variables. Consequently, this study clarified the species composition and examined correlations between spatial tree distribution and environmental factors (topographic and soil properties) in a 3 ha permanent plot in Doi Suthep-Pui National Park, Northern Thailand. The results of this research should support natural resources management, especially the selection of trees for restoration programs in mountain ecosystems.

2. METHODOLOGY

2.1 Study area

The study area in Doi Suthep-Pui National Park is located at latitude 18° 43'3" - 19° 7'30" N and longitude 98° 48'20" - 98° 57'10" E in Chiang Mai Province, Thailand (Figure 1). About 85% of the land cover is forest and 15% consists of agricultural land and urban areas (Charnsungnern and Tantanasarit, 2017). The area is characterized by a subtropical climate, with a long dry season alternating with a short, wet season; the annual precipitation is 1,139 mm and mean temperature is 24°C. The altitude ranges from 312 to 1,656 m. a. s. l. Geologically, the area contains steep hills and valleys consisting of high-grade metamorphic complex rocks and sandstone (Rhodes et al., 2005). The soil texture varies along altitudinal gradients from sandy soil and sandy clay loam at lower elevations to mostly sandy clay loam and clay soil at higher elevations (above 1,000 m. a. s. l.) (Marod et al., 2014). Complex environmental factors are found in the Doi Suthep-Pui mountain ecosystem, especially elevation gradients, which influence the distribution of forest types; these can be classified into deciduous forest (mixed deciduous forest and deciduous dipterocarp forest), mostly found in low elevations <1,000 m. a. s. l. and evergreen forest (lower montane forest and pine forest), mostly found at higher elevations (≥1,000 m. a. s. l.) (Marod et al., 2014).

2.2 Plot establishment and investigation

In 2016, a 3 ha permanent transect (50 × 600 m²) was established in the FE between DDF and LMF (latitude 18°47'4"N, longitude 98°54'58"E), where the elevation ranged from 900 to 1,100 m. a. s. l. (Figure 1). According to Hermhuk (2014) and Marod et al. (2015), three forest zones were identified based on stand clustering and canonical correspondence analysis (CCA) related to their elevation: a DDF zone from 900-940 m. a. s. l., a FE zone from 941-980 m. a. s. l. and a LMF zone above 980 m. a. s. l. The transect plot was divided into 300 subplots (10 × 10 m²). Trees with a diameter at breast height (DBH) larger than 1 cm were tagged; then, DBH was measured, positions were recorded as (x, y) coordinates, and the species were identified based on Smitinand (2014).




Figure 1. The location of the study site and permanent plot in Doi Suthep-Pui National Park, Chiang Mai Province, Thailand.

2.3 Environmental data collection

2.3.1 Topographic factors

The topographic factors (elevation, slope and aspect) of the subplots were characterized using a digital elevation model (DEM) developed by the Geo-Informatics and Space Technology Development Agency (GISTDA), Bangkok, Thailand, with a raster surface of 30 m resolution.

2.3.2 Soil properties

Samples were collected from the surface soil 0-15 cm depths in 20×50 m² subplots in the 3 ha permanent plot. In each subplot, 36 samples were collected for soil property analysis. The physical and chemical properties of the soil were analyzed at the Laboratory of Soil Science, Faculty of Agriculture, Kasetsart University, using standard methods (National Soil Survey Center, 1996). The soil texture (percentages of sand, silt and clay), soil moisture content (SMC), soil pH and amount of organic matter were determined to examine the trees' spatial distribution as a function of these properties.

2.4 Data analysis

2.4.1 Forest structure and species composition

The forest structure and species composition are described in terms of the number of individual stems, the basal area (BA) and the dominant species (Berhanu et al., 2017). The dominant species were defined using the relative frequency (RF), relative dominance (RDo), relative density (RD) and importance value index (IVI), which were estimated following Whittaker (1975) for all tree species with $DBH \geq 4.5$ cm.

2.4.2 Environmental factors

Topographic factors in this study were derived by resampling the DEM to 10 m resolution using the bilinear interpolation technique to create a more continuous surface corresponding to the spatial resolution of the field data. Estimates of topographic factors were exported for each sample. The mean elevation of a subplot was defined as the mean of the elevations of its four corners (Tarboton, 1997) and the steepest slope was the mean angular deviation from

the horizontal for each of the four triangular planes formed by connecting three of the corners of a subplot (Harms et al., 2001). Aspect in degrees relative to north was calculated using the formula:

$$\text{Aspect} = 180 - \arctan(f_y/f_x) \times (180/3.14) + 90(f_x/|f_x|),$$

where f_x and f_y are the changes in elevation when going from east to west and from north to south, respectively.

The kriging interpolation technique was used for mapping of the topographic factors and soil properties. All maps were analyzed using ArcMap ver. 10.1 (Jirakajohnkool, 2009; Walther and Meier, 2017).

The spatial distribution of tree species was considered to identify the top 20 dominant species based on IVI, combining deciduous and evergreen species. A GLM technique was applied to determine the relationships of environmental factors with tree species distribution. The dependent variable in each plot was the total basal area of each species, and the independent variables were the environmental factors. Independent variables, selected by calculating Pearson's correlation coefficients ($r \geq 0.80$ or $r \leq -0.80$) (Lan et al., 2011; Dormann et al., 2013), included nine factors: elevation, slope, aspect, soil texture (%sand, %silt and %clay), soil pH, organic matter and soil moisture content. The model omitted three factors, %silt, organic matter (SOM) and soil moisture content (SMC), because these factors were highly dependent on other selected environmental features. Consequently, six factors were used for modeling the tree spatial distributions. The model with the lowest Akaike information criterion (AIC) was chosen using the MASS package (Ripley et al., 2017) available in R version 3.2.2 according to the equation.

$$Y(I) = \text{glm}(\text{total BA}(\text{species } I) \sim x_1 + x_2 + \dots + x_6, \text{family}=\text{gaussian}(\text{link}=\text{identity})),$$

where x_1 - x_6 are independent variables: x_1 is elevation, x_2 is aspect, x_3 is slope, x_4 is %sand, x_5 is %clay and x_6 is soil pH.

3. RESULTS AND DISCUSSION

3.1 Forest structure and species composition

Within the 3 ha permanent plot, there were 11,384 tagged trees ($\text{DBH} \geq 1$ cm) across the FE, including DDF and LMF, representing 212 species,

140 genera and 67 families. Considering the trees with $\text{DBH} \geq 4.5$ cm, there were 165 species, 118 genera and 59 families. The tree density was 1,190 individual/ha and the basal area was 29.56 m²/ha. The dominant families based on species number were Fagaceae, Phyllanthaceae and Fabaceae, with 15, 12 and 10 species, respectively. These dominant families also had high total BA (m²), particularly Fagaceae and Dipterocarpaceae (Figure 2 (A)). The Fagaceae, Rubiaceae and Lauraceae had high total stem densities in the FE zone (Figure 2 (B)). *Quercus brandisiana* (32.34%) was the dominant species (based on IVI), followed by *Dipterocarpus obtusifolius* (17.73%), *Wendlandia paniculata* (17.34%), *Litsea martabanica* (17.05%), *Schima wallichii* (15.86%), *Lithocarpus garrettianus* (12.20%), *Castanopsis acuminatissima* (11.40%), *Castanopsis tribuloides* (8.90%), *Shorea obtusa* (7.96%), *Dipterocarpus tuberculatus* (6.96%), *Myrsine seguinii* (6.88%), *Tarennoidea wallichii* (6.54%), *Syzygium claviflorum* (5.24%), *Rothmannia sootepensis* (5.10%), *Persea gamblei* (5.08%), *Aporosa nigricans* (4.61%), *Lithocarpus dealbatus* (4.56%), *Scleropyrum pentandrum* (4.32%), *Aporosa villosa* (4.14%) and *Anneslea fragrans* (3.88%). These include both deciduous and evergreen species, with some species being distributed in all three zones, i.e., the DDF, FE and LMF zones, such as *Quercus brandisiana*, *Dipterocarpus obtusifolius*, *Schima wallichii*, *Wendlandia paniculata*, *Litsea martabanica* and *Anneslea fragrans* (Table 1). This indicates a high degree of success in their establishment. The species in Fagaceae, the dominant family, showed a different distribution across DDF to LMF and could be divided into two groups. The first group, consisting of *Castanopsis acuminatissima*, *Castanopsis tribuloides* and *Lithocarpus garrettianus*, was distributed mostly in the montane forest, with cooler climate conditions (Hara et al., 2002; Khamyong et al., 2003; Marod et al., 2014). The second group, consisting of *Quercus brandisiana* and *Lithocarpus dealbatus*, was found in the deciduous forest, particularly the DDF undergoing high drought conditions (Hermhuk, 2014; Marod et al., 2015). By contrast, the deciduous species in the Dipterocarpaceae (*Dipterocarpus tuberculatus* and *Shorea obtusa*) mostly occupied the DDF, with only *Dipterocarpus obtusifolius* found at elevations higher than 940 m. a. s. l., where it coexisted with evergreen species, albeit only in a narrow zone where environmental factors were similar to those of the FE (Ohsawa, 1993; Ashton, 2003).

Table 1. GLM analysis of the relationships between species distribution and environmental factors for the top 20 dominant species in the permanent plot. The values in the various columns are model regression coefficients. The coefficients with the lowest AIC were selected.

Species	Family	Type	Zonation	Environmental factors				Soil pH	
				Elevation	Aspect	Slope	%Sand		%Clay
<i>Dipterocarpus obtusifolius</i>	Dipterocarpaceae	D	All zones	-0.01***			2.16**	-1.56**	
<i>Lithocarpus dealbatus</i>	Fagaceae	D	DDF-FE	-1.24**		-1.61*		-3.37***	-1.22*
<i>Aporosa nigricans</i>	Phyllanthaceae	D	DDF-FE	-1.04*	2.23			-5.35*	-9.69**
<i>Aporosa villosa</i>	Phyllanthaceae	D	DDF-FE	-1.94**		5.57**	4.67*	-1.11*	
<i>Dipterocarpus tuberculatus</i>	Dipterocarpaceae	D	DDF	-0.01***		-0.01***	0.01**	-3.48**	-0.11**
<i>Litsea martabarnica</i>	Lauraceae	E	All zones	2.32***	5.74**	2.54**	-1.72***	2.44***	0.02*
<i>Castanopsis acuminatissima</i>	Fagaceae	E	LMF	1.96***				1.21***	
<i>Persea gamblei</i>	Lauraceae	E	LMF-FE	1.46**			-1.28**	2.01*	2.01**
<i>Scleropyrum pentandrum</i>	Santalaceae	E	LMF-FE	3.04*			-1.56*		
<i>Syzygium claviflorum</i>	Myrtaceae	E	LMF-FE	0.01***		0.01*	0.02*		
<i>Myrsine seguinii</i>	Primulaceae	E	All zones				-1.41**	1.71**	
<i>Lithocarpus garrettianus</i>	Fagaceae	E	LMF-FE		-0.9***			0.01***	-0.01*
<i>Schima wallichii</i>	Theaceae	E	All zones			3.11***		1.10***	
<i>Tarennoidea wallichii</i>	Rubiaceae	E	All zones		-7.82**	1.91**		1.21*	2.15***
<i>Castanopsis tribuloides</i>	Fagaceae	E	LMF			1.48***	-4.98**	0.15**	
<i>Anneslea fragrans</i>	Pentaphylacaceae	E	All zones			0.01*	0.15*		
<i>Wendlandia paniculata</i>	Rubiaceae	D	All zones		0.04**		0.01*		-0.07*
<i>Quercus brandisiana</i>	Fagaceae	D	All zones			0.01**	0.01**	-0.02**	
<i>Shorea obtusa</i>	Dipterocarpaceae	D	DDF-FE				3.81***	-1.25**	
<i>Rothmannia sootepensis</i>	Rubiaceae	D	DDF-FE			2.43**			

Type: D=deciduous species; E=evergreen species

Zone: LMF=lower montane forest; DDF=deciduous dipterocarp forest; FE=forest ecotone between DDF and LMF

*=p<0.05, **=p<0.01, ***=p<0.001




Figure 2. Basal area (A) and stem density (B) of each family across the DDF and LMF in Doi Suthep-Pui National Park.

3.2 Relationships of environmental factors and spatial distribution of the dominant tree species

The soil properties, especially soil texture, varied among the zones in the 3 ha permanent plot. There was a lowland-to-upland gradient in soil texture ranging from sandy to clayey types. Sandy soil was mostly distributed in the DDF and clayey soil mostly in the LMF, with a sandy clayey soil or mixed soil texture found in the FE. The soil was acidic, with a pH of 4.5-6.2 and SOM of 3.5-10% and increasing with elevation. In the FE, in addition to high pH and SOM, the SMC was high, suggesting that the observed highest tree density is due to these factors.

The spatial distribution of the 20 selected tree species reflected environmental factors, particularly elevation, although soil texture and other soil properties also influenced the distributions. The tree spatial distribution could be divided into three groups based on elevation, i.e., species found at low, intermediate and high altitudes. The distributions of the first and second groups were significantly correlated with elevation (10 species), although the correlations differed between the groups and were negative for some factors and positive for others (Table 1). By contrast, the distribution of the third group of 10 species was not significantly correlated with elevation, although slope factors strongly

influenced its distribution. The details of the spatial distributions of each group are discussed below.

The distribution of the first group was negatively associated with elevation, comprising low-altitude species, including five deciduous species: *Dipterocarpus obtusifolius*, *Dipterocarpus tuberculatus* (Figure 3(A)-(B)), *Aporosa nigricans*, *Aporosa villosa* and *Lithocarpus dealbatus*. The distribution was positively associated with sandy soil texture and acidic soil ($\text{pH} \leq 4.5$) and negatively associated with clay soil (Table 1). These findings were consistent with the species' distribution in the DDF, with its fertile soils (Teejuntuk, 2003; Sahunalu, 2009; Ueda et al., 2017). This species group can shift to higher elevation as drought or low SMC increase with climate change (Marod et al., 2002; Sahunalu, 2009; Trisurat et al., 2009).

The distribution of the second group increased with higher elevation, where the soil was mostly clayey. This group of high-altitude species included five evergreen species from the LMF: *Castanopsis acuminatissima*, *Litsea martabarnica* (Figure 3(C)-(D)), *Persea gamblei*, *Scleropyrum pentandrum* and *Syzygium claviflorum* (Table 1). In addition, distributions of *Persea gamblei* and *Litsea martabarnica* showed negative correlations with sandy soil, indicating that the families Fagaceae

(*Castanopsis acuminatissima*) and Lauraceae (*Litsea martabarnica* and *Persea gamblei*) prefer clay soil, which has good fertility based on its high organic content, as reported previously (Khamyong et al., 2003; Santisuk, 2003; Sri-Ngernyuang et al., 2003). Clay soil texture is very important for LMF

species distribution and is a common characteristic of topical montane forest (Williams-Linera, 1993; Gradstein et al., 2008; Marod et al., 2014). *Castanopsis acuminatissima* and *Syzygium claviflorum* distributions showed positive association with steep slopes (Table 1).




Figure 3. Spatial distributions of the dominant tree species in the 3 ha permanent plot as a function of elevation. The low-altitude species group includes (A) *Dipterocarpus tuberculatus* and (B) *Dipterocarpus obtusifolius*. The high-altitude species group includes (C) *Castanopsis acuminatissima* and (D) *Litsea martabarnica*.

The third group's distribution showed no significant correlation with elevation and consisted of intermediate-altitude species. Most were found in the FE and their distribution was highly dependent on soil texture and steep slopes. The group included 10 species (Table 1), with coexisting species from DDF and LMF. The distribution of the DDF species (*Quercus brandisiana*, *Shorea obtusa* (Figure 4(A)-(B)) and *Wendlandia paniculata*) were highly significantly associated with sandy soil texture (Table 1), whereas those of the LMF species (*Lithocarpus garrettianus*, *Schima wallichii* (Figure 4(C)-(D)), *Myrsine seguinii*, *Castanopsis tribuloides* and *Tarennoidea wallichii*) were associated with clayey soil. The LMF species *Schima wallichii* and *Anneslea fragrans* were distributed in all three zones (DDF, FE and LMF), indicating that they can establish in areas of drought and less fertile soil. These are light-demanding species (Asanok et al., 2013; Marod et al., 2015). The DDF species *Quercus brandisiana* and *Shorea obtusa* also had wider distributions, ranging from DDF to LMF, indicating that they have high potential to invade and establish at higher elevations when drought conditions increase and are suitable for

their regeneration (Gardner et al., 2000; Sungpalee, 2002; Santisuk, 2003; Rueangruea, 2009). By contrast, the distribution of the DDF species *Rothmannia sootepensis* was significantly correlated only with steep slopes and mostly occurred in the FE and DDF. Compared with the other species, *Rothmannia sootepensis* has larger fruit, 4.5-5 cm (Puff et al., 2005) and steep slopes may assist in seed dispersal based on gravity (Foster and Janson, 1985).

4. CONCLUSIONS

High tree species diversity was found in the FE between DDF and LMF in Doi Suthep-Pui National Park. The spatial distributions varied among species, and their ecological niches were based on environmental factors. Elevation strongly influenced the tree spatial distribution. Using elevation, the dominant species could be divided into three groups: low-altitude species (deciduous species); high-altitude species (evergreen species); and forest ecotone species (coexisting DDF and LMF species). Soil texture, which ranged from sandy to clayey, also strongly influenced on tree density. The GLM analysis also revealed that soil texture strongly

influenced the tree spatial distribution. The relationship between plant establishment and environmental factors is very important for decision

makers who establish restoration management plans. The selection of suitable species based on their niches is needed to promote successful restoration programs.




Figure 4. Spatial distribution of the intermediate-altitude species group in the 3 ha permanent plot as a function of soil texture. The distribution of the deciduous species was associated with sandy soil, including (A) *Shorea obtusa* and (B) *Quercus brandisiana*. The distribution of the evergreen group species was associated with clay texture, including (C) *Lithocarpus garrettianus* and (D) *Schima wallichii*.

ACKNOWLEDGEMENTS

This research was supported by the Center for Advanced Studies in Tropical Natural Resources (CASTNaR), Kasetsart University, Bangkok, Thailand and the Kasetsart University Research and Development Institute (KURDI). We thank the students from the Thai Forest Ecological Research Network (T-FERN) who contributed to the field observations.

REFERENCES

- Ameztegui A, Coll L. Tree dynamics and co-existence in the montane-sub-alpine ecotone: The role of different light-induced strategies. *Journal of Vegetation Science* 2011;22(6):1049-61.
- Asanok L, Marod D, Duengkae P, Pranmongkol U, Kurokawa H, Aiba M, Katabuchi M, Nakashizuka T. Relationships between functional traits and the ability of forest tree species to reestablish in secondary forest and enrichment plantations in the uplands of northern Thailand. *Forest Ecology and Management* 2013;296:9-23.
- Asanok L, Kamyo T, Norsangsri M, Salinla-um P, Rodrungruang K, Karnasuta N, Navakam S, Pattanakiat S, Marod D, Duengkae P, Kutintara U. Vegetation community and factors that affect the woody species composition of riparian forests growing in an urbanizing landscape along the Chao Phraya River, central Thailand. *Urban Forestry and Urban Greening* 2017;28:138-49.
- Ashton PS. Floristic zonation of tree communities on wet tropical mountains revisited. *Perspectives in Plant Ecology, Evolution and Systematics* 2003;6(1-2):87-104.
- Austin MP. Spatial prediction of species distribution: An interface between ecological theory and statistical modelling. *Ecological Modelling* 2002;157(2-3):101-18.
- Bacaro G, Rocchini D, Bonini I, Marignani M, Maccherini S, Chiarucci A. The role of regional and local scale predictors for plant species richness in Mediterranean forests. *Plant Biosystems* 2008;142(3):630-42.
- Berhanu A, Demissew S, Woldu Z, Didita M. Woody species composition and structure of Kuandisha afro-montane forest fragment in northwestern Ethiopia. *Journal of Forestry Research* 2017;28(2):343-55.
- Bunyavejchewin S, Lafrankie JV, Baker PJ, Kanzaki M, Ashton PS, Yamakura T. Spatial distribution patterns of the dominant canopy dipterocarp species in a seasonal dry evergreen forest in Western Thailand. *Forest Ecology and Management* 2003;175:87-101.
- Caillaud D, Crofoot MC, Scarpino SV, Jansen PA, Garzon-Lopez CX, Winkelhagen AJ, Bohlman SA, Walsh PD. Modeling the spatial distribution and fruiting pattern of a key tree species in a neotropical forest: Methodology and potential applications. *PLoS One* 2010;5(11):1-10.
- Campos JB, Romagnolo MB, Souza MCD. Structure, composition and spatial distribution of tree species in a

- remnant of the semideciduous seasonal alluvial forest of the upper Paraná River floodplain. *Brazilian Archives of Biology and Technology* 2000;43(2):185-94.
- Charnsungnern M, Tantanararit S. Environmental sustainability of highland agricultural land use patterns for Mae Raem and Mae Sa watersheds, Chiang Mai province. *Kasetsart Journal of Social Sciences* 2017;38:169-74.
- Condit R, Ashton PS, Manokaran N, LaFrankie JV, Hubbell SP, Foster RB. Dynamics of the forest communities at Pasoh and Barro Colorado: Comparing two 50-ha plots. *Philosophical Transactions of the Royal Society B: Biological Sciences* 1999;354(1391):1739-48.
- Dallmeier F. Long-term Monitoring of Biological Diversity in Tropical Forest Areas: Methods for Establishment and Inventory of Permanent Plots. Paris, France: MAB Digest 11; 1992.
- Dormann CF, Elith J, Bacher S, Buchman C, Carl G, Carré G, Marquéz JRG, Gruber B, Lafourcade B, Leitão PD. Collinearity: A review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 2013;36:27-46.
- Foster S, Janson CH. The relationship between seed size and establishment conditions in tropical woody plants. *Ecology* 1985;66(3):773-80.
- Frelich LE. Forest Dynamics and Disturbance Regimes: Studies from Temperate Evergreen-Deciduous Forests. Cambridge: Cambridge University Press; 2002.
- Fukushima M, Kanzaki M, Hara M, Ohkubo T, Preechapanya P, Choocharoen C. Secondary forest succession after the cessation of swidden cultivation in the montane forest area in Northern Thailand. *Forest Ecology and Management* 2008;255(5):1994-2006.
- Gardner S, Sidisunthorn P, Anusarnsunthorn V. A Field Guide to Forest Trees of Northern Thailand. Bangkok: Kobfai Publishing Project; 2000.
- Guisan A, Weiss SB, Weiss AD. GLM versus CCA spatial modeling of plant species distribution. *Plant Ecology* 1999;143(1):107-22.
- Guisan A, Zimmermann NE. Predictive habitat distribution models in ecology. *Ecological Modelling* 2000; 135:143-86.
- Gradstein SR, Homeier J, Gansert D. The Tropical Mountain Forest; Patterns and Processes in a Biodiversity Hotspot. Nedderrassen: Göttingen Centre for Biodiversity and Ecology; 2008.
- Hara M, Kanzaki M, Mizuno T, Noguchi H, Sri-Ngernyuang K, Teejuntuk S, Sungpalee C, Ohkubo C. The floristic composition of tropical montane forest in Doi Inthanon National Park, North Thailand, with special reference to its phytogeographical relation with montane forests in Tropical Asia. *Natural History Research* 2002;7:1-17.
- Harms KE, Condit R, Hubbell SP, Foster RB. Habitat associations of trees and shrubs in a 50-ha Neotropical forest plot. *Journal of Ecology* 2001;89(6):947-59.
- Hermhuk S. Establishment of Plant Species along the Ecotone of Lower Montane Evergreen Forest at Doi Suthep-Pui National Park, Chiang Mai Province [dissertation]. Surat Thani: Kasetsart University; 2014.
- Höfle R, Dullinger S, Essl F. Different factors affect the local distribution, persistence and spread of alien tree species in floodplain forests. *Basic and Applied Ecology* 2014;15(5):426-34.
- Homeier J, Breckle SW, Günter S, Rollenbeck RT, Leuschner C. Tree diversity, forest structure and productivity along altitudinal and topographical gradients in a species-rich Ecuadorian montane rain forest. *Biotropica* 2010;42(2):140-8.
- Jirakajohnkool S. ArcGIS 10.1 for Desktop. Bangkok: Thammasat University; 2009.
- Khamyong S, Lykke AM, Seramethakun D, Barfod AS. Species composition and vegetation structure of an upper montane forest at the summit of Mt. Doi Inthanon, Thailand. *Nordic Journal of Botany* 2003;23(1):83-97.
- Lan G, Hu Y, Cao M, Zhu H. Topography related spatial distribution of dominant tree species in a tropical seasonal rain forest in China. *Forest Ecology and Management* 2011; 262(8):1507-13.
- Li L, Huang Z, Ye W, Cao H, Wei S, Wang Z, Lian J, Sun IF, Ma K, He, F. Spatial distributions of tree species in a subtropical forest of China. *Oikos* 2009;118(4):495-502.
- Marchand P, Houle G. Spatial patterns of plant species richness along a forest edge: What are their determinants? *Forest Ecology and Management* 2006;223:113-24.
- Marod D, Kutintara U, Tanaka H, Nakashizuka T. The effects of drought and fire on seed and seedling dynamics in a tropical seasonal forest in Thailand. *Plant Ecology* 2002;161(1):41-57.
- Marod D, Sangkaew S, Panmongkal A, Jingjai A. Influences of environmental factors on tree distribution of lower montane evergreen forest at Doi Sutep-Pui National park, Chiang Mai Province. *Thai Journal of Forestry* 2014;33(3):23-33.
- Marod D, Thimkhomphang S, Panmhongkol A, Hermhuk S. Tree distribution across the forest ecotone of lower montane forest at Doi Sutep - Pui National Park, Chiang Mai Province. *Thai Journal of Forestry* 2015;34(3):99-108.
- Martin HP, Sherman RE, Fahey TJ. Tropical montane forest ecotone: Climate gradients, natural disturbance and vegetation zonation in the Cordillera Central, Dominican Republic. *Journal of Biogeography* 2007;34(10):1792-806.

- National Soil Survey Center. Soil Survey Laboratory Methods Manual. Washington: Natural Resources Conservation Service; 1996.
- Odum EP. Fundamentals of Ecology. Philadelphia: Saunders; 1959.
- Ohsawa M. Latitudinal pattern of mountain vegetation zonation in southern and eastern Asia. *Journal of Vegetation Science* 1993;4(1):13-8.
- Puff C, Chayamarit K, Chamchumroon V. Rubiaceae of Thailand: A Pictorial Guide to Indigenous and Cultivated Genera. Bangkok: Prachachon; 2005.
- Rhodes BP, Conejo R, Benchawan T, Titus S, Lawson R. Palaeocurrents and provenance of the Mae Rim Formation, Northern Thailand: Implications for tectonic evolution of the Chiang Mai basin. *Journal of the Geological Society* 2005;162(1):51-63.
- Ripley B, Venables B, Bates DM, Hornik K, Gebhardt A, Firth D, Ripley MB. Package 'mass'. CRAN Repository; 2017.
- Rueangruea S. Vegetation Structure of Montane Forest in Thailand [dissertation]. Bangkok: Kasetsart University; 2009.
- Ruiz D, Moreno HA, Gutierrez ME, Zapata PA. Changing climate and endangered high mountain ecosystem in Colombia. *Science of the Total Environment* 2008;398(1):122-32.
- Sahunalu P. Spatial distribution and size structure patterns of tree species in the long-term dynamic plots of Sakaerat deciduous dipterocarp forest, northeastern Thailand. *Journal of Forest Management* 2009;3(6):16-25.
- Santisuk T. An Account of the Vegetation of Northern Thailand. Bangkok: Royal Forest Department; 2003.
- Smitinand, T. Vegetation and Ground Covers of Thailand. Bangkok: Royal Forest Department; 2014.
- Sri-Ngernyuang K, Kanzaki M, Mizuno T, Noguchi H, Teejuntuk S, Sungpalee C, Hara M, Yamakura T, Sahunalu P, Dhanmanonda P, Bunyavejchewin S. Habitat differentiation of Lauraceae species in a tropical lower montane forest in northern Thailand. *Ecological Research* 2003;18(1):1-14.
- Sungpalee W. Some Ecological Characteristics of Deciduous Forests along the Altitudinal Gradients in Doi Inthanon National Park [dissertation]. Bangkok: Kasetsart University; 2002.
- Tang CQ, Ohsawa M. Zonal transition of evergreen, deciduous, and coniferous forests along the altitudinal gradient on a humid subtropical mountain, Mt. Emei, Sichuan, China. *Plant Ecology* 1997;133(1):63-78.
- Tarboton DG. A new method for the determination of flow directions and upslope areas in grid digital elevation models. *Water Resources Research* 1997;33(2):309-19.
- Teejuntuk S. Soil and Plant Relationships along an Altitudinal Gradient in Doi Inthanon National Park, Northern Thailand [dissertation]. Bangkok: Kasetsart University; 2003.
- Trisurat Y, Alkemade R, Arets E. Projecting forest tree distributions and adaptation to climate change in Northern Thailand. *Journal of Ecology and Natural Environment* 2009;1(3): 55-63.
- Turner IM. The Ecological of Tree in The Tropical Rain Forest. Cambridge: Cambridge University Press; 2001.
- Ueda MU, Kachina P, Marod D, Nakashizuka T, Kurokawa H. Soil properties and gross nitrogen dynamics in old growth and secondary forest in four types of tropical forest in Thailand. *Forest Ecology and Management* 2017;398:130-9.
- Valencia R, Foster RB, Villa G, Condit R, Svenning JC, Hernández C, Romoleroux K, Losos E, Magard E, Balslev H. Tree species distributions and local habitat variation in the Amazon: Large forest plot in eastern Ecuador. *Journal of Ecology* 2004;92(2):214-29.
- Walthert L, Meier ES. Tree species distribution in temperate forests is more influenced by soil than by climate. *Ecology and Evolution* 2017;7(22):9473-84.
- Weill A, Munkholm LJ. Describing soil structures, rooting and biological activity and recognizing tillage effects, damage and recovery in clayey and sandy soils. In: Ball BC, Munkholm LJ, editors. *Visual Soil Evaluation: Realizing Potential Crop Production with Minimum Environmental Impact*. Tarxien: Gutenberg Press; 2015. p. 1-13.
- Whittaker RH. *Communities and Ecosystems*. 2nd ed. New York: McMil Publicaion; 1975.
- Williams-Linera G. Soil seed banks in four lower montane forests of Mexico. *Journal of Tropical Ecology* 1993;9(3):321-37.