

Supplementary data

Population Structure and Spatial Distribution of Tree Species in Lower Montane Forest, Doi Suthep-Pui National Park, Northern Thailand

Table S1. List of species, DBH ≥ 5.0 cm, existed in 16-ha plot of LMF at Doi Suthep-pui National Park. Density (D, stems/ha), basal area (BA, m²/ha), relative basal area (RBA, %), DBH class form (DF), and important value index (IVI, %) are shown. The distribution pattern (DP) of all trees were classified into their stages, saplings (DBH < 5 cm), poling (5 $\leq$ DBH < 10 cm), and tree (DBH ≥ 10 cm).

No	Scientific name	Family	D	BA	RBA	DF	DP			IVI
							Sapling	Poling	Tree	
1	<i>Saurauia roxburghii</i> Wall.	ACTINIDIACEAE	17.69	0.10	0.31	NE	*	*	*	4.30
2	<i>Viburnum sambucinum</i> Reinw. ex Blume	ADOXACEAE	0.38	0.02	0.06	-	-	-	-	0.16
3	<i>Alangium chinense</i>	ALANGIACEAE	5.38	0.13	0.40	NE	*	*	*	1.91
4	<i>Alangium kurzii</i> Craib	ALANGIACEAE	0.06	0.02	0.06	-	-	-	-	0.07
5	<i>Alangium salvifolium</i> (L.f.) Wangerin hexapetalum	ALANGIACEAE	0.19	0.00	0.01	-	-	-	-	0.07
6	<i>Choerospondias axillaris</i> (Roxb.) B.L. Burt. & Hill	ANACARDIACEAE	6.94	0.92	2.81	NE	*	*	***	4.75
7	<i>Rhus javanica</i> L. var. javanica	ANACARDIACEAE	2.88	0.18	0.54	NE	*	***	*	1.40
8	<i>Rhus succedanea</i> L.	ANACARDIACEAE	2.19	0.03	0.09	NE	-	-	-	0.71
9	<i>Semecarpus albens</i> Kurz	ANACARDIACEAE	7.94	0.42	1.27	NE	*	***	***	3.44
10	<i>Polyalthia</i> sp.	ANNONACEAE	0.44	0.01	0.02	-	-	-	-	0.14
11	<i>Polyalthia viridis</i> Craib	ANNONACEAE	0.06	0.00	0.00	-	-	-	-	0.02
12	<i>Alstonia rupestris</i> Kerr	APOCYNACEAE	0.25	0.01	0.03	-	-	-	-	0.11
13	<i>Ilex englishii</i> Lace	AQUIFOLIACEAE	0.75	0.01	0.03	NE	*	*	*	0.26
14	<i>Ilex umbellulata</i> Loes.	AQUIFOLIACEAE	0.31	0.02	0.06	-	-	-	-	0.14
15	<i>Brassaiopsis glomerulata</i> (Blume) Regel	ARALIACEAE	0.56	0.00	0.01	NE	-	-	-	0.18
16	<i>Schefflera bengalensis</i> Gamble var. impolita Craib	ARALIACEAE	14.88	0.23	0.71	NE	*	*	*	4.41
17	<i>Schefflera elliptica</i> Harms	ARALIACEAE	2.56	0.04	0.13	NE	-	-	-	0.82
18	<i>Vernonia volkameriolia</i> Wall. ex DC.	ASTERACEAE	36.19	0.18	0.55	NE	*	*	*	7.46
19	<i>Betula alnoides</i> Buch.-Ham. ex G.Don	BETULACEAE	1.13	0.14	0.42	PO	-	-	-	0.69
20	<i>Carpinus viminea</i> Wall. ex Lindl.	BETULACEAE	0.38	0.01	0.03	-	-	-	-	0.14
21	<i>Fernandoa adenophylla</i> (Wall. ex G.Don) Steenis	BIGNONIACEAE	1.38	0.08	0.25	NE	-	-	-	0.63
22	<i>Markhamia stipulata</i> Seem.	BIGNONIACEAE	0.69	0.20	0.60	NE	***	*	*	1.36
23	<i>Stereospermum cylindricum</i> Pierre ex Dop.	BIGNONIACEAE	0.88	0.10	0.30	-	-	-	-	0.56
24	<i>Stereospermum neuranthum</i> Kurz	BIGNONIACEAE	1.75	0.03	0.10	NE	-	-	-	0.63

Table S1. List of species, DBH ≥ 5.0 cm, existed in 16-ha plot of LMF at Doi Suthep-pui National Park. Density (D, stems/ha), basal area (BA, m²/ha), relative basal area (RBA, %), DBH class form (DF), and important value index (IVI, %) are shown. The distribution pattern (DP) of all trees were classified into their stages, saplings (DBH <5 cm), poling ($5 \leq \text{DBH} < 10$ cm), and tree (DBH ≥ 10 cm) (cont.).

No	Scientific name	Family	D	BA	RBA	DF	DP			IVI
							Sapling	Poling	Tree	
25	<i>Canarium euphyllum</i> Kurz	BURSERACEAE	5.94	0.22	0.67	NE	***	***	***	2.34
26	<i>Protium serratum</i> (Wall. ex Cobbr.) Engl.	BURSERACEAE	1.00	0.02	0.07	-	-	-	-	0.34
27	<i>Glyptopetalum sclerocarpum</i> M. A. Lawson	CELASTRACEAE	0.25	0.00	0.01	-	-	-	-	0.09
28	<i>Mastixia pentandra</i> Blume subsp. <i>Chinensis</i> (Merr.) Matt.	CORNACEAE	1.06	0.02	0.05	-	-	-	-	0.31
29	<i>Diospyros glandulosa</i> Lace	EBENACEAE	4.31	0.09	0.26	NE	*	*	*	1.48
30	<i>Diospyros kerrii</i> Craib	EBENACEAE	0.88	0.02	0.06	NE	-	-	-	0.32
31	<i>Elaeocarpus floribundus</i> Blume	ELAEOCARPACEAE	4.38	0.23	0.69	NE	*	*	*	1.97
32	<i>Elaeocarpus lanceifolius</i> Roxb.	ELAEOCARPACEAE	0.38	0.03	0.10	-	-	-	-	0.21
33	<i>Elaeocarpus serratus</i> L.	ELAEOCARPACEAE	3.38	0.12	0.36	NE	*	*	***	1.32
34	<i>Sloanea sigun</i> (Blume) K.Schum.	ELAEOCARPACEAE	2.06	0.04	0.12	NE	*	*	*	0.72
35	<i>Vaccinium sprengelii</i> (G.Don) Sleumer	ERICACEAE	9.44	0.13	0.41	NE	*	*	*	2.61
36	<i>Antidesma buniis</i> (L.) Spreng.	EUPHORBIACEAE	2.25	0.05	0.16	NE	-	-	-	0.82
37	<i>Aporosa octandra</i> (Buch. -Ham ex D.Don) Vickery	EUPHORBIACEAE	16.88	0.17	0.53	NE	*	*	*	4.47
38	<i>Aporosa planchoniana</i> Baill. ex Muell. Arg.	EUPHORBIACEAE	0.25	0.00	0.01	-	-	-	-	0.08
39	<i>Baccaurea ramiflora</i> Lour.	EUPHORBIACEAE	3.38	0.04	0.12	NE	*	*	*	1.09
40	<i>Balakata baccata</i> (Roxb.) Esser	EUPHORBIACEAE	0.56	0.06	0.20	-	-	-	-	0.37
41	<i>Bischofia javanica</i> Blume	EUPHORBIACEAE	1.44	0.09	0.29	NE	-	-	-	0.68
42	<i>Bridelia glauca</i> Blume	EUPHORBIACEAE	14.31	0.31	0.96	NE	***	*	*	4.88
43	<i>Glochidion eriocarpum</i> Champ.	EUPHORBIACEAE	1.75	0.00	0.01	NE	*	*	-	0.51
44	<i>Glochidion lanceolarium</i> (Roxb.) Voigt	EUPHORBIACEAE	0.13	0.00	0.00	-	-	-	-	0.04
45	<i>Glochidion sphaerogynum</i> (Mull.Arg.) Kurz	EUPHORBIACEAE	1.56	0.01	0.03	NE	*	*	*	0.45
46	<i>Macaranga gigantea</i> (Rchb.f. & Zol.) Muell. Arg.	EUPHORBIACEAE	2.31	0.00	0.01	-	-	-	-	1.37
47	<i>Macaranga indica</i> Wight	EUPHORBIACEAE	10.56	0.54	1.63	NE	*	***	*	3.38
48	<i>Macaranga kurzii</i> (Kuntze) Pax & K. Hoffm.	EUPHORBIACEAE	0.19	0.05	0.14	NE	-	-	-	0.30
49	<i>Mallotus khasianus</i> Hook.f.	EUPHORBIACEAE	2.56	0.01	0.02	-	-	-	-	0.41
50	<i>Mallotus paniculatus</i> Mull. Arg.	EUPHORBIACEAE	1.13	0.03	0.11	NE	-	-	-	0.68
51	<i>Mallotus philippensis</i> Mull. Arg.	EUPHORBIACEAE	7.75	0.01	0.04	NE	*	*	*	1.18
52	<i>Ostodes paniculata</i> Blume	EUPHORBIACEAE	1.56	0.00	0.01	-	-	-	-	0.25

Table S1. List of species, DBH ≥ 5.0 cm, existed in 16-ha plot of LMF at Doi Suthep-pui National Park. Density (D, stems/ha), basal area (BA, m²/ha), relative basal area (RBA, %), DBH class form (DF), and important value index (IVI, %) are shown. The distribution pattern (DP) of all trees were classified into their stages, saplings (DBH <5 cm), poling ($5 \leq \text{DBH} < 10$ cm), and tree (DBH ≥ 10 cm) (cont.).

No	Scientific name	Family	D	BA	RBA	DF	DP			IVI
							Sapling	Poling	Tree	
53	<i>Phyllanthus amarus</i> Schum. & Thonn.	EUPHORBIACEAE	2.50	0.01	0.03	NE	*	*	*	0.61
54	<i>Suregada multiflorum</i> (A.Juss.) Baill.	EUPHORBIACEAE	0.06	0.00	0.00	-	-	-	-	0.02
55	<i>Symplocos macrophylla</i> Wall. ex DC.	EUPHORBIACEAE	11.31	0.10	0.30	NE	*	*	*	3.25
56	<i>Adenanthera pavonina</i> L.	FABACEAE	0.06	0.04	0.13	-	-	-	-	0.15
57	<i>Albizia chinensis</i> (Osbeck) Merr.	FABACEAE	0.06	0.00	0.00	-	-	-	-	0.02
58	<i>Albizia odoratissima</i> (L.f.) Benth.	FABACEAE	0.06	0.00	0.00	-	-	-	-	0.02
59	<i>Archidendron clypearia</i> (Jack) I.C. Nielsen	FABACEAE	5.63	0.04	0.13	NE	*	*	*	1.71
60	<i>Dalbergia assamica</i> Benth.	FABACEAE	0.81	0.03	0.09	-	-	-	-	0.30
61	<i>Dalbergia ovata</i> Graham	FABACEAE	6.13	0.14	0.44	NE	*	*	*	1.98
62	<i>Erythrina subumbrans</i> (Hassk.) Merr.	FABACEAE	2.25	0.10	0.30	NE	-	-	-	0.90
63	<i>Senna timorensis</i> (DC.) Irwin & Barneby	FABACEAE	0.06	0.00	0.00	-	-	-	-	0.02
64	<i>Castanopsis acuminatissima</i> (Blume) A.DC.	FAGACEAE	106.56	4.75	14.49	NE	*	*	*	33.60
65	<i>Castanopsis argyrophylla</i> King ex Hook.f.	FAGACEAE	0.94	0.15	0.47	NE	-	-	-	0.73
66	<i>Castanopsis armata</i> spach	FAGACEAE	29.00	1.76	5.36	NE	*	*	*	12.86
67	<i>Castanopsis diversifolia</i> (Kurz) King et Hook.f.	FAGACEAE	1.63	0.12	0.37	NE	-	-	-	0.80
68	<i>Castanopsis echinocarpa</i> Miq.	FAGACEAE	2.06	0.14	0.43	NE	-	-	-	1.06
69	<i>Castanopsis lanceifolia</i> (Oerst.) Hickel & A. Camus	FAGACEAE	0.44	0.00	0.00	-	-	-	-	0.07
70	<i>Castanopsis tribuloides</i> (Sm.) A.DC.	FAGACEAE	15.56	1.57	4.80	NE	*	***	***	8.85
71	<i>Lithocarpus annamensis</i> A. Camus	FAGACEAE	2.19	0.07	0.21	NE	*	*	*	0.84
72	<i>Lithocarpus auriculatus</i> Barnett	FAGACEAE	4.06	0.17	0.53	NE	*	*	*	1.46
73	<i>Lithocarpus dealbatus</i> (Hook.f. & Thomson) Rehder	FAGACEAE	2.94	0.07	0.20	NE	***	*	*	1.01
74	<i>Lithocarpus garrettianus</i> (Craib) A. Camus	FAGACEAE	0.06	0.00	0.01	-	-	-	-	0.03
75	<i>Lithocarpus harmandii</i> (Hick & A. Camus) A. Camus	FAGACEAE	0.25	0.03	0.09	-	-	-	-	0.15
76	<i>Lithocarpus mekongensis</i> (A. Camus) C. C. Huang	FAGACEAE	2.81	0.12	0.35	NE	*	*	***	1.11
77	<i>Lithocarpus truncatus</i> (King) Rehder & Wilson	FAGACEAE	13.69	0.67	2.03	NE	*	*	*	5.52
78	<i>Lithocarpus xylocarpus</i> (Kurz) Markgr	FAGACEAE	0.06	0.00	0.01	-	-	-	-	0.03
79	<i>Quercus oidocarpa</i> Korth	FAGACEAE	3.69	0.08	0.25	NE	*	*	*	1.21
80	<i>Casearia grewiaefolia</i> Vent. var. <i>gelonioides</i> (Blume)	FLACOURTIACEAE	0.56	0.00	0.00	NE	-	-	-	0.18

Table S1. List of species, DBH ≥ 5.0 cm, existed in 16-ha plot of LMF at Doi Suthep-pui National Park. Density (D, stems/ha), basal area (BA, m²/ha), relative basal area (RBA, %), DBH class form (DF), and important value index (IVI, %) are shown. The distribution pattern (DP) of all trees were classified into their stages, saplings (DBH <5 cm), poling ($5 \leq \text{DBH} < 10$ cm), and tree (DBH ≥ 10 cm) (cont.).

No	Scientific name	Family	D	BA	RBA	DF	DP			IVI
							Sapling	Poling	Tree	
81	<i>Flacourtia indica</i> (Burm.f.) Merr.	FLACOURTIACEAE	0.06	0.00	0.00	-	-	-	-	0.02
82	<i>Polyosma integrifolia</i> Blume	GROSSULARIACEAE	0.31	0.00	0.00	-	-	-	-	0.09
83	<i>Calophyllum polyanthum</i> Wall. ex Choisy	GUTTIFERAE	1.63	0.03	0.09	NE	-	-	-	0.58
84	<i>Garcinia cowa</i> Roxb. ex DC.	GUTTIFERAE	0.63	0.01	0.02	NE	-	-	-	0.21
85	<i>Garcinia dulcis</i> (Roxb.) Kurz	GUTTIFERAE	0.25	0.00	0.01	-	-	-	-	0.09
86	<i>Garcinia merguensis</i> Wight	GUTTIFERAE	0.56	0.00	0.01	-	-	-	-	0.17
87	<i>Garcinia speciosa</i> Wall.	GUTTIFERAE	0.69	0.01	0.02	-	-	-	-	0.23
88	<i>Garcinia thorelii</i> Pierre	GUTTIFERAE	4.25	0.09	0.27	NE	*	*	*	1.53
89	<i>Apodytes dimidiata</i> E. May. ex Arn.	ICACINACEAE	7.25	0.15	0.45	NE	*	*	*	2.53
90	<i>Engelhardtia serrata</i> Blume	JUGLANDACEAE	0.81	0.03	0.08	-	-	-	-	0.31
91	<i>Engelhardtia spicata</i> Blume	JUGLANDACEAE	4.81	0.20	0.60	NE	*	*	*	1.97
92	<i>Clerodendrum godefroyi</i>	LABIATAE	0.06	0.00	0.00	NE	-	-	-	0.02
93	<i>Vitex quinata</i> (Lour.) F.N.Williams	LABIATAE	10.44	0.41	1.26	NE	*	*	*	3.97
94	<i>Actinodaphne henryi</i> Gamble	LAURACEAE	1.25	0.04	0.13	NE	-	-	-	0.50
95	<i>Beilschmiedia gammieana</i> King ex Hook.f.	LAURACEAE	1.88	0.10	0.31	NE	-	-	-	0.87
96	<i>Cinnamomum iners</i> Reinw. ex Blume	LAURACEAE	3.06	0.08	0.24	NE	*	*	*	0.97
97	<i>Cinnamomum mollissimum</i> Hook. f.	LAURACEAE	1.63	0.03	0.09	NE	*	*	***	0.47
98	<i>Cinnamomum tamala</i> (Hamilton) Nees & Eberm.	LAURACEAE	0.88	0.10	0.29	NE	-	-	-	0.55
99	<i>Cryptocarya amygdalina</i> Nees	LAURACEAE	3.19	0.26	0.79	NE	-	-	-	1.59
100	<i>Dehaasia candolleana</i> (Meisn.) Kosterm.	LAURACEAE	0.06	0.00	0.01	-	-	-	-	0.03
101	<i>Dehaasia kurzii</i> King ex Hook. f.	LAURACEAE	1.44	0.15	0.46	NE	-	-	-	0.87
102	<i>Lindera caudata</i> (Wall. ex nees) Bth.	LAURACEAE	3.00	0.04	0.12	NE	-	-	-	0.97
103	<i>Litsea beusekomii</i> Kosterm.	LAURACEAE	1.56	0.36	1.11	NE	-	-	-	1.48
104	<i>Litsea glutinosa</i> (Lour.) C.B.Rob.	LAURACEAE	0.25	0.00	0.01	-	-	-	-	0.08
105	<i>Litsea grandis</i> (Wall. ex Nees) Hook.f.	LAURACEAE	11.50	1.03	3.15	NE	*	*	*	6.11
106	<i>Litsea martabanica</i> (Kurz) Hook.f.	LAURACEAE	27.31	0.23	0.71	NE	*	*	*	7.45
107	<i>Litsea monopetala</i> (Roxb.) Pers.	LAURACEAE	2.69	0.18	0.54	NE	-	-	-	1.22
108	<i>Litsea pierrei</i> Lecomte	LAURACEAE	2.69	0.41	1.26	NE	-	-	-	2.04

Table S1. List of species, DBH ≥ 5.0 cm, existed in 16-ha plot of LMF at Doi Suthep-pui National Park. Density (D, stems/ha), basal area (BA, m²/ha), relative basal area (RBA, %), DBH class form (DF), and important value index (IVI, %) are shown. The distribution pattern (DP) of all trees were classified into their stages, saplings (DBH <5 cm), poling ($5 \leq \text{DBH} < 10$ cm), and tree (DBH ≥ 10 cm) (cont.).

No	Scientific name	Family	D	BA	RBA	DF	DP			IVI
							Sapling	Poling	Tree	
109	<i>Neocinnamomum caudatum</i> Kosterm.	LAURACEAE	0.25	0.01	0.04	NE	-	-	-	0.22
110	<i>Neolitsea cuipala</i> (D.Don) Kosterm.	LAURACEAE	2.44	0.06	0.19	NE	-	-	-	0.66
111	<i>Persea gamblei</i> (Hook.f.) Kosterm.	LAURACEAE	24.81	0.73	2.24	NE	*	*	***	8.29
112	<i>Phoebe paniculata</i> (Nees) Nees	LAURACEAE	3.81	0.36	1.11	NE	*	*	*	2.20
113	<i>Magnolia liliifera</i> (L.) Bail.var. obovata (Korth.) Govaete	MAGNOLIACEAE	0.81	0.02	0.05	-	-	-	-	0.17
114	<i>Manglietia garrettii</i> Craib	MAGNOLIACEAE	4.50	2.10	6.41	NE	***	*	*	8.23
115	<i>Michelia baillonii</i> (Pierre) Finet & Gagnep.	MAGNOLIACEAE	1.81	0.68	2.09	NE	*	**	***	3.77
116	<i>Pterospermum grandiflorum</i> Craib	MALVACEAE	1.56	0.09	0.29	NE	*	*	*	0.76
117	<i>Osbeckia stellata</i> Buch.-Ham. ex Ker Gawl.	MELASTOMATACEAE	0.19	0.00	0.00	-	-	-	-	0.04
118	<i>Memecylon plebejum</i> Kurz	MELASTOMYRTSCEAE	0.81	0.02	0.06	NE	*	*	*	0.73
119	<i>Aglaia grandis</i> Korth. ex Miq.	MELIACEAE	1.81	0.02	0.05	NE	-	-	-	0.58
120	<i>Aglaia lawii</i> (Wight) C.J. Saldanha ex Ramamoorthy	MELIACEAE	1.13	0.09	0.26	NE	-	-	-	0.59
121	<i>Aglaia spectabilis</i> Jain & Bennet	MELIACEAE	0.44	0.00	0.01	-	-	-	-	0.14
122	<i>Heynea trijuca</i> Roxb. ex Sims	MELIACEAE	1.75	0.15	0.45	NE	*	*	*	0.95
123	<i>Melia azedarach</i> L.	MELIACEAE	3.81	0.09	0.27	-	-	-	-	0.87
124	<i>Toona ciliata</i> M. Roem.	MELIACEAE	0.25	0.04	0.12	-	-	-	-	0.20
125	<i>Antiaris toxicaria</i> Lesch.	MORACEAE	0.38	0.01	0.03	-	-	-	-	0.13
126	<i>Artocarpus chaplasha</i> Roxb.	MORACEAE	0.44	0.04	0.12	-	-	-	-	0.26
127	<i>Artocarpus nitidus</i> Trecul	MORACEAE	0.75	0.01	0.02	-	-	-	-	0.25
128	<i>Artocarpus rigidus</i> Blume	MORACEAE	0.19	0.00	0.00	-	-	-	-	0.06
129	<i>Ficus auriculata</i> Lour.	MORACEAE	0.44	0.01	0.02	-	-	-	-	0.15
130	<i>Ficus curtipes</i> Corner	MORACEAE	0.63	0.00	0.02	NE	-	-	-	0.18
131	<i>Ficus hirta</i> Vahl	MORACEAE	0.06	0.00	0.01	-	-	-	-	0.02
132	<i>Ficus racemosa</i> L.	MORACEAE	0.06	0.00	0.00	-	-	-	-	0.02
133	<i>Ficus semicordata</i> Buch.-Ham. ex Sm.	MORACEAE	0.44	0.00	0.01	NE	-	-	-	0.14
134	<i>Ficus subincisa</i> Sm.	MORACEAE	0.06	0.00	0.00	-	-	-	-	0.02
135	<i>Morus macroura</i> Miq.	MORACEAE	0.81	0.23	0.71	NE	*	*	*	1.35
136	<i>Horsfieldia glabra</i> (Blume) Warb.	MYRISTICACEAE	0.31	0.00	0.01	-	-	-	-	0.11

Table S1. List of species, DBH ≥ 5.0 cm, existed in 16-ha plot of LMF at Doi Suthep-pui National Park. Density (D, stems/ha), basal area (BA, m²/ha), relative basal area (RBA, %), DBH class form (DF), and important value index (IVI, %) are shown. The distribution pattern (DP) of all trees were classified into their stages, saplings (DBH <5 cm), poling ($5 \leq \text{DBH} < 10$ cm), and tree (DBH ≥ 10 cm) (cont.).

No	Scientific name	Family	D	BA	RBA	DF	DP			IVI
							Sapling	Poling	Tree	
137	<i>Horsfieldia ridleyana</i> (King.) Warb.	MYRISTICACEAE	0.13	0.00	0.00	-	-	-	-	0.04
138	<i>Ardisia corymbifera</i> Mez	MYRSINACEAE	0.13	0.00	0.01	-	-	-	-	0.05
139	<i>Ardisia quinqueгона</i> Blume	MYRSINACEAE	0.44	0.01	0.03	-	-	-	-	0.16
140	<i>Maesa ramentacea</i> (Roxb.) A.DC.	MYRSINACEAE	0.19	0.00	0.01	NE	*	*	*	0.19
141	<i>Rapanea yunnanensis</i> Mez	MYRSINACEAE	3.00	0.08	0.25	NE	*	*	*	1.10
142	<i>Decaspermum parviflorum</i> (Lam.) A.J.Scott	MYRTACEAE	0.63	0.01	0.02	-	-	-	-	0.21
143	<i>Syzygium claviflorum</i> (Roxb.) a.M.Cowan & Cowan	MYRTACEAE	5.25	0.27	0.83	NE	*	*	*	2.18
144	<i>Syzygium helferi</i> (Duthie) Chantar. & J.Parn.	MYRTACEAE	4.44	0.20	0.62	NE	-	-	-	1.67
145	<i>Syzygium tetragonum</i> (Wight) Walp.	MYRTACEAE	6.75	0.94	2.87	NE	*	**	*	4.77
146	<i>Syzygium toddlioides</i> (Wight) Walp.	MYRTACEAE	11.25	0.95	2.89	NE	****	*	*	6.21
147	<i>Chionanthus ramiflorus</i> Roxb	OLEACEAE	1.13	0.01	0.02	NE	-	-	-	0.32
148	<i>Fraxinus floribunda</i> Wall. ex Roxb.	OLEACEAE	0.56	0.10	0.29	-	-	-	-	0.45
149	<i>Olea rosea</i> Craib	OLEACEAE	0.88	0.02	0.06	NE	*	*	*	0.54
150	<i>Olea salicifolia</i> Wall. ex G.Don	OLEACEAE	0.06	0.01	0.03	NE	-	-	-	0.21
151	<i>Pittosporum napaulense</i> (DC.) Rehder & E. H. Wilson	PITTOSPRACEAE	0.13	0.01	0.02	-	-	-	-	0.06
152	<i>Dacrycarpus imbricatus</i> (Blume) de Laub.	PODOCARPACEAE	0.88	0.01	0.03	-	-	-	-	0.25
153	<i>Podocarpus neriifolius</i> D.Don	PODOCARPACEAE	0.69	0.03	0.09	PO	-	-	-	0.28
154	<i>Helicia nilagirica</i> Bedd.	PROTEACEAE	22.44	0.29	0.89	NE	*	*	*	6.32
155	<i>Heliciopsis terminalis</i> (Kurz) Sleumer	PROTEACEAE	1.13	0.04	0.12	NE	-	-	-	0.46
156	<i>Carallia brachiata</i> (Lour.) Merr.	RHIZOPHORACEAE	2.88	0.10	0.31	NE	*	*	*	1.16
157	<i>Eriobotrya bengalensis</i> (Roxb.) Hook. f.	ROSACEAE	5.50	0.08	0.26	NE	*	*	*	1.74
158	<i>Prunus arborea</i> (Blume) Kalkman	ROSACEAE	3.25	0.04	0.11	NE	***	*	*	1.04
159	<i>Prunus cerasoides</i> D.Don	ROSACEAE	0.25	0.00	0.01	-	-	-	-	0.08
160	<i>Prunus ceylanica</i> (Wight) Miq.	ROSACEAE	0.06	0.02	0.07	-	-	-	-	0.09
161	<i>Canthium glabrum</i> Blume	RUBIACEAE	1.13	0.02	0.05	NE	-	-	-	0.39
162	<i>Oxyceros bispinosus</i> (Griff.) Tirveng.	RUBIACEAE	0.06	0.00	0.00	-	-	-	-	0.02
163	<i>Pavetta indica</i> L.	RUBIACEAE	1.44	0.02	0.05	NE	-	-	-	0.42
164	<i>Psydrax nitida</i> (Craib) K.M.Wong	RUBIACEAE	0.25	0.01	0.02	-	-	-	-	0.09

Table S1. List of species, DBH ≥ 5.0 cm, existed in 16-ha plot of LMF at Doi Suthep-pui National Park. Density (D, stems/ha), basal area (BA, m²/ha), relative basal area (RBA, %), DBH class form (DF), and important value index (IVI, %) are shown. The distribution pattern (DP) of all trees were classified into their stages, saplings (DBH <5 cm), poling ($5 \leq \text{DBH} < 10$ cm), and tree (DBH ≥ 10 cm) (cont.).

No	Scientific name	Family	D	BA	RBA	DF	DP			IVI
							Sapling	Poling	Tree	
165	<i>Randia pilosa</i> Craib	RUBIACEAE	0.38	0.00	0.00	-	-	-	-	0.06
166	<i>Tarennoidea wallichii</i> (Hook.f.) Tirveng. & Sastre	RUBIACEAE	3.25	0.17	0.53	NE	***	*	***	1.50
167	<i>Wendlandia paniculata</i> (Roxb.) DC.	RUBIACEAE	15.69	0.16	0.47	NE	*	*	*	4.22
168	<i>Acronychia pedunculata</i> (L.) Miq.	RUTACEAE	13.25	0.40	1.23	NE	*	*	*	4.66
169	<i>Clausena excavata</i> Burm. f.	RUTACEAE	0.19	0.00	0.00	-	-	-	-	0.05
170	<i>Clusiana lansium</i> (Lour.) Skeels	RUTACEAE	2.31	0.01	0.03	NE	*	*	*	0.70
171	<i>Glycosmis pentaphylla</i> (Retz.) DC.	RUTACEAE	0.44	0.00	0.00	NE	-	-	-	0.09
172	<i>Micromelum minutum</i> (G.Forst.) Wight & Arn.	RUTACEAE	4.00	0.00	0.01	NE	-	-	-	0.62
173	<i>Meliosma pinnata</i> Walp.	SABIACEAE	8.69	0.00	0.01	-	-	-	-	1.19
174	<i>Sarcosperma arboreum</i> Hook.f.	SACOSPERMATACEAE	5.25	0.24	0.74	NE	*	*	*	2.19
175	<i>Scleropyrum wallichianum</i> (Wight & Arn.) Arn.	SANTALACEAE	0.63	0.01	0.02	-	-	-	-	0.18
176	<i>Mischocarpus pentapetalus</i> (Roxb.) Radlk.	SAPINDACEAE	0.69	0.03	0.08	NE	-	-	-	0.37
177	<i>Nephelium hypoleucum</i> Kurz	SAPINDACEAE	0.06	0.01	0.05	-	-	-	-	0.12
178	<i>Madhuca floribunda</i> (Pierre) H.J.Lam	SAPOTACEAE	0.94	0.00	0.00	-	-	-	-	0.13
179	<i>Picrasma javanica</i> Blume	SIMAROUBACEAE	1.63	0.03	0.10	NE	-	-	-	0.58
180	<i>Duabanga grandiflora</i> (Roxb. ex DC.) Walp.	SONNERATIACEAE	0.44	0.15	0.45	-	-	-	-	0.58
181	<i>Turpinia pomifera</i> (Roxb.) DC.	STAPHYLEACEAE	20.06	0.32	0.98	NE	*	*	*	5.97
182	<i>Sterculia lanceolata</i> Cav.	STERCULIACEAE	0.19	0.00	0.01	-	-	-	-	0.06
183	<i>Styrax benzoides</i> Craib	STYRACACEAE	38.94	0.34	1.03	NE	*	*	*	9.95
184	<i>Symplocos cochinchinensis</i> (Lour.) S.Moor	SYMPLOCACAEAE	1.06	0.03	0.10	NE	-	-	-	0.44
185	<i>Adinandra integerrima</i> T. Anderson ex Dyer	THEACEAE	1.06	0.05	0.16	NE	-	-	-	0.46
186	<i>Anneslea fragrans</i> Wall.	THEACEAE	0.38	0.00	0.00	-	-	-	-	0.12
187	<i>Camellia oleifera</i> Abel	THEACEAE	0.94	0.02	0.06	NE	-	-	-	0.34
188	<i>Camellia pleurocarpa</i>	THEACEAE	1.63	0.03	0.08	NE	-	-	-	0.57
189	<i>Eurya acuminata</i> DC.	THEACEAE	7.63	0.08	0.25	NE	*	*	*	2.29
190	<i>Pyrenaria diospyricarpa</i> Kurz	THEACEAE	7.06	0.15	0.44	NE	*	*	*	2.23
191	<i>Schima wallichii</i> (DC.) Korth.	THEACEAE	19.13	3.03	9.23	NE	*	***	*	14.61

Table S1. List of species, DBH ≥ 5.0 cm, existed in 16-ha plot of LMF at Doi Suthep-pui National Park. Density (D, stems/ha), basal area (BA, m²/ha), relative basal area (RBA, %), DBH class form (DF), and important value index (IVI, %) are shown. The distribution pattern (DP) of all trees were classified into their stages, saplings (DBH <5 cm), poling ($5 \leq \text{DBH} < 10$ cm), and tree (DBH ≥ 10 cm) (cont.).

No	Scientific name	Family	D	BA	RBA	DF	DP			IVI
							Sapling	Poling	Tree	
192	<i>Ternstroemia gymnanthera</i> (Wight & Arn.) Bedd.	THEACEAE	0.69	0.02	0.06	NE	-	-	-	0.25
193	<i>Trema angustifloia</i> (Planch.) Blume	ULMACEAE	0.31	0.00	0.01	-	-	-	-	0.09
194	<i>Xanthophyllum siamensis</i> Craib	XANTHOPHYLLACEAE	0.06	0.00	0.00	-	-	-	-	0.02
195	<i>Xanthophyllum virens</i> Roxb.	XANTHOPHYLLACEAE	3.31	0.13	0.41	NE	*	*	*	1.37
			806.88	32.79	100	122	72	72	72	300